

FEEDING COWS ON VEGETABLES AND FRUITS MARKET WASTES AND ITS EFFECT ON CHEMICAL COMPOSITION OF MILK

Mousa, M. A. M.

Dairy Chemistry Laboratory, Animal Prod. Res. Inst. Sakha Exp. Station for Agric. Res. Sakha, Kafr El-Sheikh, Egypt.

ABSTRACT

Chemical composition of milk as affected by using vegetables and fruits market waste (VFMW) in feeding cows was examined over a period of 12 weeks. The use of VFMW in a silage in (Treatment I) to replace 15% of the control ration (Ammonia-treated rice straw and concentrate) resulted in an increase in fat, lactose, protein, casein and whey protein content of milk. Non-protein N and non-casein N were lower. In treatment (II), VMW and El-Mufeed was used to replace 15% of the control ration, whereas in treatment (III), VMW and ground yellow corn was used to replace 5% of the control ration. Results showed that treatment (II) gave the highest fat and lactose content followed by treatment (I), whereas treatment (I) gave the highest protein and casein content followed by treatment (II). In all cases, the control milk samples had the lowest values of fat, protein, casein and whey protein content.

Keywords: milk composition, cows feeding

INTRODUCTION

In Egypt especially in the long summer days, it is well known that farm animals suffer from an extreme shortage of green forages and supplying the animals with sufficient quantity of nutrients is the most critical problem. Only straw, maize stover and other fibrous materials are available which are considered as poor and inadequate sources for protein and minerals. Recently, search for unconventional sources for animal feeds is the target for many studies: El-Shinnawy (1974) and Tag El-Din (1979) used some by-products in a mixture for feeding farm animals. Osman (1980) used poultry waste in ruminant nutrition, Anis and El-Abd (1991) supplemented rice straw and maize stover with El-Mufeed (a molasses liquid supplement) and mentioned that such treatment increased milk yield and its nutritional value, Abo El-Nor *et al.* (1995) used date seeds as a new component in diet for dairy animals, whereas El-Feel *et al.* (1995) followed the trace elements and composition of milk as affected by feeding cows on poultry manure, Khattab *et al.* (1998) and El-Alamy *et al.* (1999) used feather and offal meals and other protein sources in feeding buffaloes.

Concerning vegetables and fruits market waste (VFMW), Abaza *et al.* (1987) and Gabr *et al.* (1995) evaluated the nutritive value of VFMW and concluded that it can participate in solving the feed shortage problem. The present work was conducted to study the effect of using VFMW in feeding the

lactating cows on gross chemical composition and nitrogen distribution of the milk.

MATERIALS AND METHODS

1- Material

Vegetables and fruits marketing wastes (VFMW) were collected from the biggest Egyptian market for vegetables and fruits (Al-Aopour market) and preserved into silage as described by Gabr *et al.* (1995). The VFMW contained a mixture of unhomogenous materials such as cabbage and cauliflower leaves, leek, lettuce, onion, radish, tomatoes, potato, peas....etc. This mixture of materials contained about 85% moisture.

- El-Mufeed is a commercial name for a molasses-based liquid supplement rich in urea, minerals and vitamins

2- Experimental animals

In the present study, 20 Friesian lactating cows weighted 400-500 Kg were divided randomly into 4 dietary treatments. Treatments were arranged to be one control and the other three for the tested rations (5 cows for each treatment). The feeding trials were lasted for 3 months.

3- Experimental rations

The cows were individually fed according to the allowances for dairy cows. The control group was fed on traditional summer ration in Egypt which was composed of 3% ammonia-treated rice straw and concentrate. In treatment (I), the ration of this group was the same as in the control with the substitution of 15% by VFMW in a silage form. In treatment (II), the ration was the same as in the control with the substitution of 15% by vegetables marketing waste (VMW) and El-Mufeed. The ration of treatment (III) was the same as the control ration with the substitution on of 5% by VMW and ground yellow corn.

4- Analytical procedures

The weekly milk samples from evening and morning milking were collected over a period of 12 weeks and chemically analysed for fat and lactose using Milko-scan (133 B. Foss Electric), whereas the total protein content was determined using semi-micro Kjeldahl method as given by LING (1963). The N distribution was done as recommended by Rowland (1938). Non-casein nitrogen (NCN) was determined in the resultant filtrate after precipitation of casein by means of acetic acid and sodium acetate at pH 4.6. Non-protein nitrogen (NPN) was measured in the clear filtrate after precipitation of protein by TCA (15%). Casein nitrogen (CN) and whey protein nitrogen (WPN) were quantified by difference specified by Rowland (1938) as follows:

$$\text{CN} = \text{Total N} - \text{NCN.}$$

$$\text{WPN} = \text{NCN} - \text{NPN.}$$

Statistical methods

The obtained data were statistically analysed for range, average and standard error (SE) as well as for the analysis of variance using a SAS computer program as give in the user's guide (SAS, 1987).

RESULTS AND DISCUSSION

Table 1 shows chemical composition of milk as affected by the feeding treatments applied. Fat content ranged from 1.09 to 3.42% in control milk, whereas the corresponding values 1.91-4.90%, 1.44-4.43% and 1.57-4.78% for treatment I, II, III, respectively. The maximum average fat content of 2.99±0.07% was given for treatment (II) followed by milk from treatment (I), whereas the minimum average value of 2.48±0.08% was recorded for the control milk.

Table (1): Effect of feeding treatments on the chemical composition (%) of milk *.

Property	Treatments			
	Control	I	II	III
Fat content				
Range	1.09-3.42	1.91-4.90	1.44-4.43	1.57-4.78
AV±SE	2.48±0.08	2.94±0.07	2.99±0.07	2.82±0.08
Lactose content				
Range	4.18-5.37	4.20-5.37	4.69-5.25	4.17-5.08
AV±SE	4.69±0.04	4.87±0.03	4.91±0.02	4.64±0.03
Protein content				
Range	1.85-4.72	1.98-4.15	2.55-4.59	1.85-5.23
AV±SE	3.02±0.07	3.36±0.06	3.28±0.05	3.23±0.09
Casein content				
Range	1.08-2.93	1.21-3.38	1.79-3.13	1.08-3.89
AV±SE	2.14±0.07	2.52±0.06	2.45±0.04	2.41±0.08
Whey protein content				
Range	0.06-1.15	0.25-0.83	0.13-0.89	0.19-1.08
AV±SE	0.45±0.03	0.50±0.04	0.51±0.04	0.51±0.04

* Range, average and standard error (SE) of 60 determinations.

The increase in milk fat content in milk from different treatments compared to the control might be due to the higher energy and the increased digestible crude protein for the supplemented rations. Anis *et al.* (1991) observed the same trend of results when they used EI-Mufeed in feeding lactating cows.

Lactose content (Table 1) shows the same trend of fat with the exception that the lowest average lactose content was recorded for treatment (III). However, lactose content ranged from 4.18 to 5.37% in control milk, whereas those for treatments I, II and III were 4.20-5.37, 4.69-5.25 and 4.17-5.08% in order. The corresponding averages ± SE were 4.69±0.04, 4.87±0.03, 4.91±0.02 and 4.64±0.03% respectively. Such variations in lactose content due to FT were highly significant and those due to FP were only significant (Table 2).

Table (2): Analysis of variance for the effect of feeding treatments (FT) and feeding period (FP) on chemical composition of milk.

SOV	DF	SS	MS	F	Significance <
Fat content					
FT	3	8.559	2.853	10.17	0.01
FP	11	7.879	0.716	2.55	0.01
FTxFP	33	15.205	0.460	1.64	0.05
Lactose content					
FT	3	2.523	0.841	14.88	0.01
FP	11	1.280	0.116	2.06	0.05
FTxFP	33	2.737	0.083	1.47	NS*
Protein content					
FT	3	3.408	1.136	6.67	0.01
FP	11	15.086	1.371	8.05	0.01
FTxFP	33	17.862	0.541	3.18	0.01
Casein content					
FT	3	6.404	2.135	12.61	0.01
FP	11	13.031	1.185	7.00	0.01
FTxFP	33	15.293	0.463	2.74	0.01
Whey protein content					
FT	3	0.169	0.056	3.06	0.05
FP	11	2.124	0.193	10.47	0.05
FTxFP	33	2.374	0.072	3.90	0.05

* NS = not significant

Protein content (Table 1) in milk from each group of cows showed a wide range of variations. Thus, in control milk, the min. and max. values were 1.85 and 4.72% in order. The corresponding values in case of treatment (I) were 1.98 and 4.15%, and in treatment (II) were 2.55 and 4.59% and in treatment (III) were 1.85 and 5.32%, in order. The maximum average value of $3.36 \pm 0.06\%$ was given for treatment (I). This was followed by values of 3.28 ± 0.05 , 3.23 ± 0.09 and $3.02 \pm 0.07\%$ for treatments II, III and the control respectively.

Statistical analysis (Table 2) showed that the differences in protein content due to FT, FP and FT x FP were highly significant.

Similar results were given by Anis *et al.* (1991) when they used El-Mufeed and this could be attributed to the effect of sufficient digestible protein and starch value of the ration to satisfy all or some requirements of lactating cows. Gabr *et al.* (1995) demonstrated that the VFMW hay contain crude protein as 11-36%, and when VFMW silage was added to maize stover at the rate of 1:1 and 2:1, it caused an increase in its crude protein from 5.20 to 8.76% and 11.66%, respectively.

The pattern of casein content was the same as that observed for protein content. Table (1) reveals that the maximum average in this respect ($2.52 \pm 0.06\%$) was recorded for treatment (I) with the range of 1.21-3.38%, whereas the minimum average ($2.14 \pm 0.07\%$) was given for the control milk with the range of 1.08-2.93%. Treatment (II) had an average of $2.45 \pm 0.04\%$ with the range of 1.79-3.13%, whereas those for treatment (III) were $2.41 \pm 0.08\%$ and 1.08-3.89% in order.

The same highly significant effect of FT, FP and FTxFP was recorded in Table (2) as that mentioned for protein content.

In spite of the same average value ($0.51 \pm 0.04\%$) was record for whey protein content of milk from treatments (II) and (III), the minimum and maximum given values were completely different. However, a wider range of variation was observed with respect to whey protein of the control milk. The values ranged from 0.06 to 1.15% with the minimum average value of $0.45 \pm 0.03\%$. However, the differences in whey protein due to FT, FP and FTxFP were significant at $P < 0.05$. As given in Table (3), TN followed the same pattern of protein content being the highest average value of 0.51% was recorded in milk from treatment (I), whereas the control milk had the minimum value of 0.47%. This was true with respect to CN. Their values had the same trend of casein content. The maximum average value of $0.40 \pm 0.02\%$ was recorded for treatment (I). This was followed by the values of 0.39 ± 0.02 , 0.36 ± 0.01 and $0.34 \pm 0.01\%$ for milk samples from treatment II, III and the control respectively.

Table (3): Effect of feeding treatments on nitrogen distribution (%) in milk *.

Property	Treatments			
	Control	I	II	III
TN content				
Range	0.31-0.58	0.31-0.65	0.41-0.65	0.29-0.82
AV±SE	0.47±0.01	0.51±0.02	0.50±0.02	0.50±0.01
CN content				
Range	0.19-0.59	0.19-0.53	0.30-0.57	0.14-0.62
AV±SE	0.34±0.01	0.40±0.02	0.39±0.02	0.36±0.01
NPN content				
Range	0.03-0.15	0.02-0.06	0.03-0.08	0.01-0.11
AV±SE	0.06±0.004	0.04±0.002	0.05±0.002	0.07±0.003
NCN content				
Range	0.09-0.21	0.08-0.16	0.08-0.19	0.11-0.23
AV±SE	0.14±0.004	0.12±0.002	0.13±0.002	0.15±0.004
WPN content				
Range	0.01-0.18	0.04-0.13	0.02-0.14	0.03-0.13
AV±SE	0.07±0.005	0.08±0.004	0.08±0.004	0.08±0.004

* Range, average and standard error (SE) of 60 determinations.

Statistical analysis (Table 4) revealed that FT, FP and FTxFP had highly significant effects on TN and CN content.

It may be of interest to point out that the maximum average values of $0.07 \pm 0.003\%$ and $0.15 \pm 0.004\%$ for NPN and NCN, in order, were recorded for milk from treatment (III), whereas the minimum values of $0.04 \pm 0.002\%$ and $0.12 \pm 0.002\%$, in order, were given for treatment (I) as shown in Table (3). However, Table (4) shows that the differences in NPN and NCN due to FT, FP and FTxFP were highly significant.

Table (4): Analysis of variance for the effect of feeding treatments (FT) and feeding period (FP) on nitrogen distribution in milk.

SOV	DF	SS	MS	F	Significance <
TN content					
FT	3	0.074	0.024	5.28	0.01
FP	11	0.352	0.032	6.85	0.01
FTxFP	33	0.452	0.013	2.93	0.01
CN content					
FT	3	0.171	0.056	12.48	0.01
FP	11	0.349	0.031	6.98	0.01
FTxFP	33	0.343	0.011	2.28	0.01
NPN content					
FT	3	0.042	0.014	14.23	0.01
FP	11	0.032	0.002	2.96	0.01
FTxFP	33	0.073	0.002	2.23	0.01
NCN content					
FT	3	0.027	0.009	27.20	0.05
FP	11	0.020	0.002	5.60	0.05
FTxFP	33	0.055	0.002	5.03	0.05
WPN content					
FT	3	0.589	0.196	1.31	NS*
FP	11	1.428	0.130	< 1	NS
FTxFP	33	4.911	0.149	< 1	NS

* NS = not significant

Table (3) reveals that the average WPN value of $0.08 \pm 0.004\%$ was recorded for milk samples from treatments I, II and III, whereas a slight lower value of $0.07 \pm 0.005\%$ was given for the control milk. Such differences in WPN as affected by FT, FP and FTxFP were insignificant. In general, such change in chemical composition of milk as affected by using the market wastes might be due to the chemical composition of the given wastes. Gabr *et al.* (1995) gave the values of 8.76, 2.18, 20.30, 44.80 and 23.96% on dry matter basis for crude protein, ether extract, crude fiber, nitrogen-free extract and ash respectively for the mixture containing VFMW and maize stover as 1:1. The corresponding values in the mixture of 2:1 was 11.66, 2.47, 21.02, 44.10 and 20.75% in order.

REFERENCES

- Abaza, M.A.; A.M. Nour; B.E. Borhami and K. El-Shazly (1987). Nutritive value of vegetable and fruit wastes as animal feeds. *Alex.J.Agric. Res.* 32, 75-83.
- Abo El-Nor, S.A.H.; H.M. El-Sayed and A.M. Kholif (1995). Date seeds as a new component in diets for dairy animals. *Egyptian J. Dairy Sci.* 23, 167-176.
- Anis, S.M.K. and M.M. El-Abd (1991). Supplementation of field by-products by EL-Mufeed in relation to milk production and composition. *Egyptian J. Dairy Sci.* 19, 145-155.
- El-Alamy, H.A.; A.M. Kholif; M.A. EL-Ashry; H.M. El-Sayed and S.M. Kholif (1999). Yield and composition of buffaloes milk and animal health as affected by including feather and offal meals in the diets. *Egyptian J. Dairy Sci.* 27, 13-25.

- El-Feel, F.M.; F.S. Ibrahim; Z.B.H. Rabie and E.A.M. Abd Alla (1995). Trace elements and milk composition of cow's milk fed on poultry manure. Egyptian J.Dairy Sci. 23: 87-99.
- El-Shinnawy, M.M. (1974). The use of some by-products in feed mixture for feeding animals. Ph.D.Thesis, Fac.of Agric., Cairo Univ., Egypt.
- Gabr,A.A.; A.Z. Mehrez; M.Y. El-Ayek and E.Kh. Hamed (1995). Nutritional evaluation of market wastes from fruits and vegetables as feeded for sheep. Proc. 5th Sci. Conf. Animal Nutrition.1: 179-189.
- Khattab, H.M.; H.A. El-Alamy; S.A.H. Abo El-Nor; F.A.F. Salem and M.M.A. Abdou (1998). Effect of protein source on milk yield and composition of lactating buffaloes. Egyptian J.Dairy Sci. 26, 1-23.
- Ling, E.R. (1963). A Text Book of Dairy Chemistry. vol.2, 3rd ed. Chapman and Hall Ltd, London.
- Osman, A.M.A. (1980). Using poultry waste in ruminant nutrition. M.Sc. Thesis, Fac.of Agric., El-Minia Univ., Egypt.
- Rowland, S.J. (1938). The determination of the nitrogen distribution in milk. J.Dairy Res. 9, 42-46.
- SAS. (1987). Statistical Analysis System. User's Guide, Version 6th ed., SAS Institute, N.C., USA.
- Tag El-Din, A.E.S. (1979). Utilization of some by-products in the feeding of ruminants. M.Sc. Thesis, Fac. of Agric.,Alex. Univ., Egypt.

تأثير استخدام مخلفات أسواق الفاكهة والخضراوات فى تغذية الأبقار على التركيب الكيماوى للبن

محمد عرفه محمد موسى

قسم بحوث كيمياء الألبان - معهد بحوث الإنتاج الحيوانى - مركز البحوث الزراعية - وزارة الزراعة

أهتم البحث بدراسة تأثير استخدام مخلفات أسواق الفاكهة والخضراوات كغذاء للأبقار على التركيب الكيماوى للبن حيث استخدمت هذه المخلفات فى صورة سيلاج لمدة ١٢ اسبوعاً وفقاً للمعاملات الآتية:

- ١- عليقة المقارنة (الكنترول) كانت عبارة عن قش معاملة بالأمونيا والعليقة المركزة.
 - ٢- فى المعاملة الأولى تم استبدال ١٥% من مخلوط عليقة الكنترول بمخلفات خضراوات وفاكهة كمصدر للطاقة.
 - ٣- فى المعاملة الثانية تم استبدال ١٥% من مخلوط عليقة الكنترول بمخلفات خضراوات مع مفيد.
 - ٤- فى المعاملة الثالثة تم استبدال ٥% من مخلوط عليقة الكنترول وخضراوات + ذرة أصفر.
- وقد دلت النتائج على أن التغذية وفقاً للمعاملة الأولى أدت لزيادة محتوى اللبن من الدهن واللاكتوز والبروتين الكلى والكازين وبروتينات الشرس بينما انخفض النتروجين غير البروتينى والنتروجين غير الكازينى. وقد لوحظ أن تأثير المعاملة الثانية كان أكبر عن المعاملة الأولى فى زيادة نسبة الدهن واللاكتوز وتأثيراً أقل على زيادة نسبة البروتين والكازين. وبصفة عامة فقد اتصف اللبن الناتج من التغذية على عليقة المقارنة بمحتويات أقل من الدهن والبروتين والكازين وبروتينات الشرس.