

UTILIZATION OF SOME FRUITS AND VEGETABLES WASTES AS A SOURCE OF DIETARY FIBERS IN CAKE MAKING

Sharoba, A.M.* ; M. A. Farrag** and A. M. A. Abd El-Salam**

* Food Sci. Dept., Moshtohor Fac. of Agric. Banha Univ., Egypt.

** Food Technology Res. Institute, Agric. Res. Center, Giza, Egypt.

ABSTRACT

In this study, feasibility of using orange waste (OW), carrot pomace (CP), potato peels (PP) and green pea peels (GPP) wastes from food industry, as a starting raw material to produce dietary fibre powders and the feasibility of producing cakes intended for people suffering from obesity or over weight and diabetes. The physicochemical properties of dietary fiber powder were first evaluated. The results showed that the fibre contents, compositions and hydration properties (water and oil holding capacity and swelling capacity) of the fiber powder. Some food processing wastes {(OW), (CP), (PP) and (GPP)}, are a rich sources of fibres. The OW, CP, PP and GPP wastes were replaced with wheat flour (72%) at 5, 10, 15 and 20% levels and studied for rheological characteristics. Water absorption increased significantly with increasing wastes from 0 to 20%. Dough stability and dough development were increased. Resistance to extension values significantly increased, whereas extensibility values decreased. The produced fiber substituted cakes achieved a reduction in calories. Cakes were prepared from blends of wheat flour (72%) with 5, 10, 15 and 20% fruits and vegetables wastes. The volume of cakes decreased with increase in fruits and vegetables wastes content from 0 to 20%. Cakes prepared from 20% of wastes OW, CP, PP and GPP had a higher ratio of dietary fibres. Sensory evaluation showed that all high fiber substituted cake samples were significantly lower than control cake sample in all sensory characteristics, except cake samples prepared with 5 and 10% of orange waste and carrot pomace had no significant differences ($P>0.05$) with control cake. However, there were significant differences between cake samples containing the same type of fiber source at 5, 10, 15 and 20% replacement levels. The highest and lowest scores in the same type of fiber source for all attributes were that achieved by cake samples with fiber source at 5 and 20% replacement levels, respectively. Texture properties of cake were affected by replacement levels of fiber source. The results indicated that OW, CP, PP and GPP wastes can serve as a good source of dietary fibres.

Keywords: Food processing wastes; dietary fibers; orange waste, carrot pomace, potato peels, green pea peels; cakes; rheological, sensory and texture characteristics.

INTRODUCTION

Dietary fiber (DF) consists of a variety of non starch polysaccharides which include cellulose, hemicellulose, pectin, B-glucans, gums and lignin (Lamghari *et al.*, 2000 and Gallaher and Schneeman, 2001). However, the SDF/IDF ratio is important for both, dietary and functional properties. It is generally accepted that those fiber sources suitable for use as food ingredient should have an SDF/IDF ratio close to 1:2 (Jaime *et al.*, 2002). Dietary fiber is composed mainly by remnants of edible plant cells; parenchymatous tissues

are known to be the most important source of fiber, cell walls of fruits, vegetables, pulses and cereals make up most of the dietary fibre intake (Jimenez *et al.*, 2000).

Dietary fiber plays an important role in human health, they acts as a protective agent against cardiovascular diseases, diverticulosis, constipation, irritable colon, colon cancer and diabetes (Gorinstein *et al.*, 2001 and Rodriguez *et al.*, 2006). Fiber derived from fruits and vegetables have a considerably higher proportion of soluble dietary fiber, whereas cereal fibres contain more insoluble cellulose and hemicellulose (Redgwell and Fischer, 2005 and Herbafood, 2002). The physiological effects are related to the physicochemical and functional properties of dietary fibre. It is widely known that dietary fibres obtained by different methods and from different sources, behave differently during their transit through the gastrointestinal tract, depending on their chemical composition and physicochemical characteristics and on the processing that food undergo (Jimenez *et al.*, 2000; Herbafood, 2002 and Chau and Huang, 2003). While a few years ago the sub products generated during the processing of plant food constituted an economic and environmental problem, today they are considered a promising source of functional compounds (Carle *et al.*, 2001).

They are many fruits, for example orange, apple and peach, which are used for the extraction of their juices. They all contain a by-product from which can be recovered different high-added value compounds; among those, it is remarkable the fibre fraction that has a great potential in the preparation of functional foods. There are also several vegetables, such as carrot, potato, green pea, pepper, artichoke, onion and asparagus that originate a waste during their processing (Rodriguez *et al.*, 1999 and Rodriguez *et al.*, 2006) and it contains both soluble and insoluble fibre compounds that can be used for designing new 'functional foods'. Orange and other fruits which are mainly commercialized in processed form originate great amounts of sub-products consisting of peels, bones and seeds. This material could be a restrictive factor in the commercialization of these products if it is not usefully recovered, because it represents significant losses with respect to the raw material, which considerably increases the price of the processed products (Schieber *et al.*, 2001).

Potato peels consider as a rich source of fibers, since it contains about 9.7 to 68% from its total weight, Mullin *et al.* (1993). The potatoes are the most commonly consumed vegetables worldwide, whereas the potatoes production stands at 293 million metric tons. The annual production of Egypt is 2.2 million tons. As results, a large quantity of potato peel waste is generated.

Green peas (*Pisum Sativum*) which is one of the most important vegetable processing crop in Egypt and it's by product (peels) resembles about 50%. Dietary fibres from different sources have been used to replace wheat flour in the preparation of bakery products. Potato peel, a by-product from potato industry, rich in dietary fibre, was used as a source of dietary fiber in bread making (Toma *et al.*, 1979).

Plant fibres show some functional properties, such us water-holding capacity (WHC) and swelling capacity (SWC) which have been more useful

for understanding the physiological effect of dietary fiber, than the chemical composition alone (Femenia *et al.*, 1997; Gallaher and Schneeman, 2001). Functional properties of plant fibre depend on the IDF/SDF ratio, particle size, extraction condition and vegetable source (Jaime *et al.*, 2002). The main characteristics of the commercialized fiber product are: total dietary fiber content above 50%, moisture lower than 9%, low content of lipids, a low caloric value and neutral flavour and taste (Larrauri *et al.*, 1997 and Larrauri 1999). In order to take advantage of the dietary and functional properties of fibre, some high dietary fibre formulated foods are currently being developed (Grigelmo-Miguel and Martin-Belloso, 1999; Herbafood, 2002; Tudorica *et al.*, 2002). According to Larrauri (1999), the "ideal dietary fibre" should meet, among others, the following requirements; have no nutritionally objectionable components, be as concentrated as possible, be bland in taste, color and odor; have a balanced composition; have a good shelf life; be compatible with food processing; have the expected physiological effects. It must be kept in mind that fiber enrichment not only influences the overall quality of food by changing its physiological properties, but also significantly affects the sensorial properties of a product. When plant fibres are added to a food product, they contribute to water holding properties and texture of the product (Kethireddipalli *et al.*, 2002). Fruit fibre have better quality due to higher total and soluble fibre content, water and oil holding capacity (Larrauri, 1999). Residues from orange juice extraction are potentially an excellent source of DF because this material is rich in pectin and may be available in large quantities (Fernandez-Gines *et al.*, 2003). Dietary fibres are not only desirable for their nutritional properties, but also for their functional and technological properties and because of those they can also be used to upgrade agricultural products and by products for use as food ingredients (Schieber *et al.*, 2001). The industry of fruit juice produces significant amounts of by-products which could cause problems in their disposal. Usually, these products are used in animal feeding. However, their high amount of dietary fiber could permit the use of them in developing new natural ingredients for the food industry. According to the Statistical Database of the Food and Agriculture Organization of the United Nations (FAOSTAT), world orange production in 2007 was estimated to be 63,906,094 tons. A high percentage of this production (70%) is used to manufacture products such as juice or marmalade. Moreover, approximately 50–60% of the processed fruit is transformed into citrus peel waste, which is composed of the peel, seeds and membrane residues (Wilkins *et al.*, 2007). In order to prevent problems related to the disposal of this product and environmental concerns, this waste must be properly processed. Under current environmental legislation, all waste must be considered raw material if a valorization procedure is to be developed.

Accordingly, the purpose of this work was a trial to evaluate some functional properties of fibre obtained from orange waste, carrot pomace, potato peels and green pea peels wastes, in order to use them as a dietary fibres source in the enrichment of cakes. The obtained cakes were evaluated mechanically, physically and organoleptically.

MATERIALS AND METHODS

Materials:

Wheat flour: Wheat flour (soft) (72%), were obtained from the South Cairo Mills Company, Cairo, Egypt.

Baking Ingredients: skim dry milk, sugar, margarine, corn oil, eggs, vanilla and baking powder were obtained from local market Kaha city, Kaliobia, Egypt.

Orange waste (*Citrus sinensis*) Balady orange variety: were obtained from Kaha company for canned food, Kaha city, Kaliobia, Egypt.

Carrot pomace (*Daucus carota* L.): was obtained from Kaha company for canned food, Kaha city, Kaliobia, Egypt.

Potato peels (*Solanum tuberosum* L.): were obtained as a waste from Egypt Foods factory for chippy products, Industrial zone, Minufiya, Egypt.

Green pea peels (*Pisum Sativum*): were obtained from Kaha Company for canned food, Kaha city, Kaliobia, Egypt.

Preparation dietary fiber powders

•**Orange waste fibres:** the by-products obtained from orange peel and the remaining pulp after juice extraction could be suitable sources of DF by cutting, extraction of juice, peel residue chopping, the material was washed under mild conditions to avoid or minimize losses of some soluble fiber components (such as pectins and pentosans) as well as bioactive components (such as flavonoids, polyphenols and carotenes) (Larrauri, 1999), then dried at temperatures below 65 °C avoids changes in the functional properties and in the content of polyphenols, tannins, anthocyanidins and proteins for 12 hrs. in an electric oven drier, and grinding to a particle size of 500–600 µm, according to (Kethireddipalli *et al.*, 2002).

•**Carrot pomace fibers:** Carrot pomace was washed twice with warm water (30 °C); then it was dried at 60 °C for 12 hrs. in an electric oven drier and ground to a particle size of 500–600 µm. to pass through 20 mesh sieve packed in polyethylene bags and stored in refrigerator until use.

•**Potato peels fiber:** potato peels obtained and thoroughly washed on a multilayer of cheese cloth using tap water and then dried in an electric oven at 60 °C for 12 hrs. to prepare dried potato peels.

•**Green pea peels fiber:** green pea peels were washed with water; then it was dried at 60 °C for 12 hrs. in an electric oven drier and ground to a particle size of 500–600 µm. to pass through 20 mesh sieve packed in polyethylene bags and stored in refrigerator until use.

The fibre materials were then finely milled using the laboratory hammer mill, and sieved through a 500-µm sieve size into fine powder. Four batches of fibre flours were prepared. The fibres were assigned names, orange waste fibre (OWF), Carrot pomace fibre (CPF), Potato peels fibre (PPF) and Green pea peels fibre (GPPF), then packaged in polythene bags protected from light and moisture and stored at refrigerator temperature until further analysed.

Processing:

Preparation of cakes

The corn oil was beaten thoroughly, the sugar was added to butter and mixed until got smooth like cream, and then a well blended egg with vanilla were added and mixed together. The blends soft wheat flour (72%) with dietary fibre sources (orange waste fibre (OWF), Carrot pomace fibre (CPF), Potato peels fibre (PPF) and Green pea peels fibre (GPPF) these by-products were replaced with wheat flour at 5, 10, 15 and 20% levels, baking powder were stirred together and added alternately to the egg mixture. The mixture was whipped until got smooth. The dough transferred to a greased pan and was baked for 25 min. at $200\pm 5^{\circ}\text{C}$ then was cooled at room temperature (AACC, 2002). Cakes were prepared according to the formula is shown in Table (I).

Table (I): Cake formula:

Ingredients	Weight (g)
Soft wheat flour (72% extraction)	150.00
Baking powder	6.81
Salt	3.40
Sugar	75.00
Corn oil	31.83
Fresh whole egg	39.75
Skim dry milk	14.76
Vanilla	1.50

Analytical methods:

Proximate chemical composition:

Moisture, crude protein, crude lipid, ash and carbohydrate contents were determined using the appropriate AOAC (2000).

Carbohydrates were determined by difference from the total dietary fibre, lipids, protein and ash contents (Chau and Huang, 2003).

Determination of dietary fiber contents:

Total dietary fiber (TDF), soluble dietary fibre (SDF) and insoluble dietary fiber (IDF) contents of samples were determined with an enzymatic–gravimetric procedure according to AOAC Method 991.43, (AOAC, 2000 and Lee *et al.*, 1992).

Functional properties:

Water holding capacity (WHC) and oil holding capacity (OHC) were measured according the methods reported by Femenia *et al.* (1997) and Robertson *et al.* (2000).

Swelling capacity (SWC): Was measured using the bed volume technique described by Kuniak and Marchessault (1972).

Triplicate measurements were taken for all WHC, OHC and SWC.

Rheological characteristics:

Fruits and vegetables waste blends at 0, 5, 10, 15 and 20% levels were prepared by replacing wheat flour. The effect of fruits and vegetables fiber on the mixing profile of the dough was studied using farinograph (Brabender, Duisburg, Germany) according to the standard AACC methods

(2000). Farinograph test was carried out to determine the water absorption, arrival time, dough development time, dough stability and degree of weakening.

The elastic properties of the dough were studied using extensograph (Brabender, Duisburg, Germany) according to the standard AACC methods (2000). Extensograph test was carried out to determine resistance to extension (B.U.), extensibility (mm), proportional number and energy (cm²).

Physical characteristics for cakes:

The weight (g) for cake was determined individually within one hour after baking the average was recorded.

The volume (cm³) of different types of produced cakes was determined by rape seeds displacement method according to (AACC, 2000). Specific volume was calculated according to the method of (AACC, 2000), using the following equation

$$\text{Specific volume} = \text{Volume (cm}^3\text{)}/\text{Weight (g)}.$$

Texture characteristics:

The texture of the cakes was measured objectively using food texturometer (TAHDi, Stable Micro System, UK) as per the standard AACC methods (2000). A test speed of 2.0 mm s⁻¹ were used. A 35 mm diameter cylinder aluminum probe (P-35), was used to measure the required compression force. Force required to compress 25% of the cake slice (2.54 cm) was recorded. All measurements were performed at ambient temperature 25±2°C according to Gomez *et al.* (2007).

Mechanical properties measurements:

All mechanical properties were made using the Instron Universal Testing Machine (Model 4401) equipped with: 5-mm diameter tip probe for penetration test (with 100, 500, 1000 and 5000 N load cell). All testing was performed at room temperature (25±2°C).

Penetration test (PT):

Each piece of tested cakes was placed in a hole of the bevelled ring. The pin penetrated with a constant speed 10 mm.min⁻¹ into each piece of cakes tested. Three penetration points at different parts for each piece of cakes and 5 piece for each sample were taken for each test condition, recording the force profile at 1 cm of probe penetration (Femenia *et al.*, 1997). Force – deformation curves were recorded and hardness was derived as indicators of textural properties. Each sample was used for only one measurement.

Mechanical measurements analysis:

1- The force corresponding to the maximum puncture is defined as the maximum force (F_{max}). The maximum puncture force (F_{max}) was measured in Newtons (N), as mentioned by Sharoba *et al.* (2012) and Saleh *et al.* (2012).

2- Maximum deformation: the distance from beginning to distance at maximum force.

3- Hardness = Maximum force (N) / Maximum deformation (mm), as mentioned by Sharoba *et al.* (2012) and Saleh *et al.* (2012).

Sensory analysis

Sensory evaluation of cakes were carried out in a standardized test room in morning sessions (11:00-13:00 h) by a 12 trained sensory panel, and

were carried out by a properly well trained panel of 12 panelists. They were selected if their individual scores in 10 different tests showed a reproducibility of 90%. The 12 member internal panel evaluated on a five point hedonic scale, scoring was based on a 100 point scale (10-100) where (90-100) = excellent, (70-80) = very good, (50-60) = good, (30-40) = fair and (10-20) = poor. Mineral water was used by the panelists to rinse the mouth between samples, according to the method described by (AACC, 2002), cake samples were left to cool at room temperature for 1 hr. after baking. Then cake was cut with a sharp knife and subjected to panel test. Cells 30 (uniformity 10, size of cells 10, and thickness of walls 10), grain 20, texture 30 (moistness 10, tenderness 10, and softness 10), crumb color 10, flavor 10 and overall acceptability 100 degrees.

Statistical analysis:

Each sample was prepared in triplicate and analysed twice. Data are presented as the average of three analyses and data was analysed using the analysis of variance (ANOVA) $p < 0.05$ was considered to be statistically significant, followed by multiple comparisons using least significant difference, (L.S.D._{0.05})(Gomez and Gomez, 1984).

RESULTS AND DISCUSSION

Nowadays, there is a considerable interest in studying the feasibility of using by-products from food processing plants as raw materials for production of dietary fiber powder since these wastes are inexpensive and highly abundant. Fruits and vegetables wastes or residues are among the most promising by-products with high dietary fiber content, and possess a good balance between SDF and IDF

Proximate chemical composition of wheat flour (72 %):

Data in Table (1) represent the proximate chemical composition of wheat flour (72 %) of protein, fat, ash, crude fiber, total carbohydrates contents on (dry weight basis) and moisture contents. From these results the wheat flour was suitable for used to make cake.

Table (1): Chemical composition of wheat flour (72 %).

Components	%
Moisture	11.99 ± 0.292
Protein*	11.85 ± 0.401
Fat*	1.06 ± 0.068
Ash*	0.52 ± 0.018
Crude fiber*	0.54 ± 0.034
Available carbohydrates**	86.04

* (on dry weight basis)

** Available carbohydrate by difference

Chemical composition, fiber content and functional properties of used fruits and vegetables wastes:

Although interest in fiber as a food ingredient has waned in recent years, the importance of fiber cannot be overlooked. The health benefits coupled with functional properties such as water and fat holding capacity

properties have created a renewed interest in fiber, particularly in the nutraceutical industry. Many of the fiber supplements which have been researched are obtained from by-products resulting from the processing of fruits and vegetables. Increased use of fiber supplementation would therefore not only improve the health benefits and functional properties of many foods. Data in Table (2) represent the proximate chemical composition of fruits and vegetables wastes, the results are in agreement with those obtained by Sosulski and Wu (1988); Camire and Flint (1991); Ralet *et al.* (1993); Camire *et al.*, (1997) and Chantaro *et al.* (2008).

Table (2): Proximate chemical composition of used fruits and vegetables wastes.

Components	dietary fibers source			
	Orange waste	Carrot pomace	Potato peels	Green pea peels
Moisture (%)	4.15±0.32	4.61±0.21	3.58±0.63	4.28±0.27
Total ash (%)	10.03±0.54	7.29±0.32	6.92±0.27	7.18±0.34
Total fat (%)	1.57±0.02	1.75±0.01	2.25±0.08	1.34±0.03
Total protein (%)	8.72±0.36	10.06±0.18	12.16±0.48	13.27±0.51
Total dietary fiber (%)	63.45	69.85	73.25	71.30
Insoluble fiber (%)	41.17±1.28	45.12±1.08	53.39±0.94	51.48±1.34
Soluble fiber (%)	22.28±0.93	24.73±1.22	19.86±1.47	19.82±1.36

Functional properties:

It is well known that the functional properties of dietary fibers have the greatest effect on their functions in foods and physiological action in the body El-Refai *et al.*, 2006. The results of the functional properties of fruits and vegetables wastes are given in Table (3). The functional properties of plant fibre depend on the IDF/SDF ratio, structure of the plant polysaccharides and vegetables or fruits sources.

Table (3): Functional properties of used fruits and vegetables wastes

Components	dietary fibers source			
	Orange waste	Carrot pomace	Potato peels	Green pea peels
Water holding capacity (WHC) (g water/g dry fiber source)	16.39±0.47	19.72±0.38	15.62±0.67	13.48±0.14
Swelling capacity (SWC)	20.74±0.62	23.96±0.58	16.27±0.38	14.93±0.41
Oil holding capacity (OHC) (g oil/g dry fiber source)	1.27±0.01	3.95±0.17	8.63±0.48	6.89±0.35

The water holding capacity (WHC) is the quantity of water that remains bound to the hydrated fiber following the application of an external force (pressure or centrifugation), It consists of the sum of bound water, hydrodynamic water and, mainly, physically trapped water (Raghavendra, *et al.*, 2006 and Vazquez- Ovando *et al.*, 2009). It is an important property of DF from both a physiological and technological point of view. Dietary fiber holds water by adsorption and absorption phenomena and some water is also retained outside the fiber matrix (free water). WHC depends on fiber

processing, washing increases WHC probably due to the removal of sugars, and the chemical and physical structure: the water-holding capacity is related to the SDF content, and high levels of SDF produce a high WHC because soluble fibres, possess a higher WHC than cellulosic fibres. WHC of different dietary fiber sources; orange waste, carrot pomace, potato peels and green pea peels are presented in Table 3. Relatively high values of WHC were obtained compared with the values obtained with other agricultural by-products including apple pomace and pear pomace (Grigelmo-Miguel and Martin-Belloso, 1999). The high WHC values were related to the soluble dietary fiber fraction and pectic substances contained in the peels might account for their high WHC in orange waste. High WHC was also previously reported for orange waste and carrot pomace. The water holding capacity of carrot pomace was higher than that of orange waste. The higher WHC for the samples under study because we used suitable drying temperature to prepare dietary fiber sources. The WHC was lower at a higher drying temperature; this might be due to the degradation of some soluble dietary fiber components, leading to the loss of ability to retain water in the sample. The values of WHC are 16.39, 19.72, 15.62 and 13.48 for orange waste, carrot pomace, potato peels and green pea peels, respectively which are higher than those reported for others fibrous residues such as grape, mango, peach and tomato (7.8-9.4 g water/g product) (El-Refai *et al.*, 2006), bambangans peel (11.6 g water/g product), apple pomace (8.4 g water/g product) (Sudha *et al.*, 2007 and Hassan *et al.*, 2011), and other dietary fiber products such as date paste (1.3 g water/g product) (Sanchez-Zapata *et al.*, 2011) and similar with other dietary fiber products such as carrot peel (21.89 g water/g product) (Chantaro *et al.*, 2008). The high WHC values could indicate a lightness improve of food products where they were added.

The SWC of carrot pomace are presented in Table (3). The results showed that DF powder from carrot pomace possessed high SWC. These values are of similar order with those of cauliflower fiber (Femenia *et al.*, 1997), coconut fiber (Raghavendra *et al.*, 2006) and carrot fiber (Chantaro *et al.*, 2008) but higher than those reported for citrus residues (Figuerola *et al.*, 2005). Blanching had a significant effect on the SWC. During blanching carrot, some components might be lost with water and the change of structural tissues might enhance the water uptake. However, the other fiber sources blanching step don't occurred. Furthermore, a very high correlation between the SWC and WHC was obtained at each dietary fiber sources. In addition to chemical compositions, some physical properties, such as structure, particle size, porosity, pH, temperature, ionic strength, types of ions in solutions and density are important to the understanding of the deferent behaviors of samples during hydration.

The two hydration properties, SWC and WHC, which are mainly determined by the food content (like dietary fiber) have been shown to be closely related, oil-holding capacity OHC is another important property of food ingredients used in formulated food. The (OHC) is also a technological property related to the chemical structure of the plant polysaccharides and depends on surface properties, overall charge density, thickness, and hydrophobic nature of the fiber particle (Figuerola *et al.*, 2005; Carme *et al.*,

2007 and Fernandez-López *et al.*, 2009); it is also related with the IDF content, particle size and drying, dehydration promotes a general decrease in fiber OHC compared with the fresh fiber. The fiber sources show an OHC of 1.27, 3.95, 8.63 and 6.89 g oil/g dry fibre respectively. This is similar the OHC of orange by-products (1.81 g oil/g fibre), lemon by-products (6.60 g oil/g fiber) and tiger nut by-product (6.90 g oil/g fibre) (Lario *et al.*, 2004; Figuerola *et al.*, 2005 and Sanchez-Zapata *et al.*, 2009). For this reason, foods added with potato peels or green pea peels will not retain high amounts of oil, both in the case that it will be added as ingredient in the food product, or in the case that it will be used for frying processes.

This suggested that the all fiber sources in this study would be able to stabilize food emulsions with a high percentage of fat. Basically, the mechanism of OHC is mainly due to the physical entrapment of oil by capillary attraction. Moreover, the hydrophobicity of proteins also plays a major role in fat absorption. Therefore, among the fiber sources samples, the variations in OHC may be partially due to the different proportions of polar side chains of the amino acids on the surfaces of their protein molecules. Furthermore, the OHC of fiber sources samples are also related to the particle size, overall charge density and hydrophilic nature of the individual particles. Similarly, the correlation between OHC and total amount of protein and TDF was very high. This implied that the OHC of the fiber sources samples might also depend on the total content of protein and TDF present.

Rheological properties:

Dough produced from blends of wheat flour (72%) and different levels of dietary fibers (orange waste, carrot pomace, potato peels and green pea peels) were tested for rheological properties.

Farinograph parameters of wheat flours dough blended with dried fibers sources:

The effect of replacing wheat flour with different levels (5, 10, 15 and 20 %) of dried dietary fiber sources (orange waste, carrot pomace, potato peels and green pea peels) on mixing dough properties are presented in Table (4). Data in Table (4) shows the effect of blending fiber sources with wheat flour (72 %) on the farinograph parameters, i.e. water absorption, arrival time, dough development time, dough stability and dough weakening.

The results indicate that water absorption, arrival time and dough development time were increased due to the increase of wheat flour replacement levels. This might be due to the increase in the fiber level in dietary fiber sources blends. These results agree with those obtained by Abd El-Moniem and Yassen (1993); who reported that addition of fiber sources caused an increase in the water absorption of the produced dough. This may be due to higher water hydration capacity of fibers. Dough stability increased due to the increase of wheat flour replacement with dietary fiber sources. These results confirmed increased stability with those obtained by Doweidar (2001); who found that the addition of dietary fiber source (dry carrot and pectin). Also, reported that the addition of dietary carrot led to increase water absorption and stability. Degree of weakening decreased due to the increase replacement levels of wheat flour with dietary fiber sources, this might be due to increasing of fibrous materials, which contribute gluten net development.

Table (4): Effect of dietary fibres sources addition at different levels to wheat flour on farinograph parameters of dough.

Dietary fiber source	Replacement levels	Farinograph parameters				
	(%)	Water absorption (%)	Arrival time (min)	Dough development (min)	Dough stability (min)	Degree of weakening (B.U.)
Control	0	61.50	1.50	2.75	6.00	95.25
Orange waste	5	67.40	1.75	3.30	6.40	80.25
	10	68.10	2.30	4.25	6.85	75.25
	15	70.90	3.00	5.50	7.25	69.75
	20	72.45	4.10	6.00	7.70	54.00
Carrot pomace	5	63.30	1.90	3.20	6.40	83.80
	10	65.15	2.30	3.90	6.90	77.50
	15	68.95	2.90	4.30	7.30	69.90
Potato peels	20	71.95	3.80	5.10	7.95	57.80
	5	63.50	1.85	3.10	6.80	83.40
	10	65.95	2.25	3.75	7.45	69.00
Green pea peels	15	67.55	2.55	4.25	8.25	57.20
	20	69.85	2.90	4.60	9.10	47.95
	5	62.75	1.70	3.00	6.75	77.10
Green pea peels	10	63.45	2.10	3.45	7.85	60.20
	15	64.75	2.25	3.80	9.40	48.65
	20	65.90	2.45	4.05	10.80	35.60

Water absorption of the control (wheat flour 72% ext.) showed a value of 61.5 ml. While, wheat flour blended with different ratios of fibre sources, 5, 10, 15 and 20% showed a gradual increase in parallel with fiber sources increasing, this may be due to the high fiber contents in fiber sources. Arrival time showed a time in min ranged between 1.30 to 4.10 min.. Dough development time of blends with fiber sources at the ratios of 5, 10, 15 and 20% resulted in (2.75 to 6 min). Dough stability showed its maximum value (10.8 min.), when the blend contained 20% dry green pea peels and 80% wheat flours 72% extraction followed by 20% dry potato peel and 80% wheat flours 72% extraction (9.1 min.). From the above mentioned data, it could be concluded that, the replacement of fiber sources at different ratios improved the gluten network and farinograph parameters of the wheat flour dough. From the aforementioned data, it could be concluded that blending with fiber sources had a good farinograph parameters.

Extensograph parameters of wheat flours dough (72% extraction) blended with dried fibre sources:

Data in Table (5) shows the effect of addition fiber sources (in ratios of 5, 10, 15 and 20%) with wheat flours (72% extraction) dough on the extensograph parameters, i.e. extensibility (E), resistance to extension (R) and the ratio between them and also the energy. The results show that the resistance was 781.25 B.U. for the control sample (wheat flour 72% ext.), while it decreased with increasing the levels of fiber sources in prepared blends. Also, the resistance value of dough produced from blends of wheat flour (72% ext.) with dry green pea peels or potato peels more than of dough produced from blends for wheat flours (72% ext.) with orange waste or carrot pomace but both still decreased than the control sample with increasing the

fiber sources ratio. Extensibility (E) showed a value of 157.5 mm for the control sample (wheat flour 72% ext.). Blending with fiber sources showed a slight decrease due to the dried fiber sources. The lowest extensibility value was recorded due to the blend containing 20% green pea peels. The ratio between resistances to extension of extensibility reflects the effect of blends on the elasticity of the produced dough. Concerning the energy, the control sample resulted in the highest value while, other treatments resulted in the following decreasing. From the above mentioned data, it could be concluded that addition of fibre sources had a good extensograph parameters like the control sample.

Table (5): Effect of dietary fibers sources addition at different levels to wheat flour on extensograms parameters of dough.

Dietary fiber source	Replacement levels	Extensograph parameters			
	(%)	Resistance to extension (B.U.)	Extensibility (m.m.)	Proportional number (R / E)	Energy (cm ²)
Control	0	781.25	157.50	4.96	127.50
Orange waste	5	790.00	133.80	5.90	113.75
	10	786.85	125.00	6.29	105.00
	15	748.95	113.80	6.58	101.50
	20	691.10	108.30	6.38	96.00
Carrot pomace	5	729.50	154.50	4.72	125.20
	10	715.35	145.05	4.93	115.80
	15	683.20	133.35	5.12	106.40
	20	675.00	125.80	5.37	99.30
Potato peels	5	795.60	129.00	6.17	119.40
	10	742.45	121.45	6.11	107.95
	15	695.50	116.35	5.98	98.65
	20	639.00	110.00	5.81	90.80
Green pea peels	5	840.05	113.50	7.40	118.95
	10	808.35	103.00	7.85	111.00
	15	783.50	90.55	8.65	103.95
	20	744.70	82.80	8.99	93.45

Effect of addition dietary fiber sources on texture characteristics of cakes substituted by different levels fiber:

The results presented in Table (6) show the texture characteristics (mechanical properties) of cake contained different types of fiber source, firmness must be explained by the different chemical interactions between oil, protein starch and fibre that affect its retrogradation, the interactions between the swollen starch granules, fibre and the protein network actively contribute to crumb firming, the results showed increased of cake firmness, with the exception of control cake sample. Cake contained the different levels of the dietary fiber sources led to the hardest with firmness higher than that of the control cake sample. Fiber can provoke a crumb firmness increase probably due to the thickening of the crumb cells surrounding the air spaces. Also firmness was increase may be a result of molecular entanglements between fiber and gluten proteins (Gomez et al., 2007).

Table (6): Effect of addition dietary fiber sources on texture characteristics of cakes substituted by different levels of fibers

Characteristics	Replacement levels (%)	Dietary fiber sources			
		Orange waste	Carrot pomace	Potato peels	Green pea peels
Firmness	0 (Control)	8.905	8.905	8.905	8.905
	5	9.736	10.331	9.792	10.277
	10	11.629	12.035	10.720	13.056
	15	13.681	14.168	12.297	15.641
	20	16.751	16.738	14.638	17.356
Cohesiveness	0 (Control)	0.683	0.683	0.683	0.683
	5	0.602	0.581	0.521	0.624
	10	0.492	0.346	0.453	0.537
	15	0.308	0.292	0.258	0.395
	20	0.239	0.207	0.171	0.311
Gumminess	0 (Control)	2.894	2.894	2.894	2.894
	5	3.132	3.028	2.947	3.354
	10	3.538	3.387	3.205	3.725
	15	3.892	3.897	3.439	3.962
	20	3.984	3.972	3.620	4.108
Chewiness	0 (Control)	2.963	2.963	2.963	2.963
	5	2.985	3.025	2.974	3.342
	10	3.129	3.195	3.047	3.649
	15	3.367	3.408	3.115	3.803
	20	3.658	3.827	3.248	3.958
Springiness	0 (Control)	0.742	0.742	0.742	0.742
	5	0.692	0.655	0.673	0.718
	10	0.583	0.525	0.534	0.634
	15	0.490	0.497	0.468	0.587
	20	0.405	0.378	0.361	0.435
Resilience	0 (Control)	0.697	0.697	0.697	0.697
	5	0.612	0.607	0.530	0.581
	10	0.542	0.510	0.401	0.421
	15	0.421	0.383	0.252	0.382
	20	0.310	0.306	0.197	0.285

An increase in chewiness and gumminess values with ageing was observed in all recipes. Chewiness is one of the texture parameters easily correlated with sensory evaluation through trained panels (Esteller *et al.*, 2004). Both, gumminess and chewiness are parameters dependant on firmness. Therefore, their values followed a similar trend than that of firmness. Cohesiveness quantifies the internal resistance of food structure. As happened with firmness, cake cohesiveness depended on the addition of fiber. A similar result was also obtained in other baked goods (Esteller *et al.*, 2004). A subjective evaluation of springiness is normally made by consumers and consists of slightly pressing the piece of food, by hand or with the mouth, and verifying how easily it returns to the original size. Finally, the addition of dietary fiber sources to wheat flour to produced cake improved the texture properties of cake due to high water holding capacity of fibre, these results

are in agree with many researchers which have shown in their studies that remarkable the lack of any clear relationship between cake water contents and its firmness. Decreased crumb firmness did not improve crumb texture, which showed increased crumbliness.

Effect of addition types of dietary fiber source on hardness measurement (mechanical properties) of cake

Data listed in Table (7) show the effect of addition different types of fiber sources with wheat flour (72%) to produced cake on compression properties (hardness measurement). The results show that, all additions were increase hardness by increasing addition levels comparing with control sample. The results in the same table showed that, the addition fiber sources to produced cakes improved the hardness due to the effect of fibers, the fibers' function had high adsorption capacity of oil and water and therefore the hardness increased with adding fibers. These results agree with Sowmya *et al.* (2009). The hardness of the cake ranged from 0.524 to 0.820 for control sample and the addition of 20% green pea peels sample, respectively. The above results indicated that the addition of types of fiber sources is beneficial in improving the maximum force and texture.

Table (7): Effect of addition dietary fiber sources on hardness measurement (mechanical properties) of cakes

Dietary fiber sources	Substituted levels	Maximum Force	Maximum deformation	Hardness
	(%)	N	mm	N/mm
Control	0	26.22	50	0.524
Orange waste	5	28.37	50	0.567
	10	30.04	50	0.601
	15	33.31	50	0.666
	20	36.94	50	0.739
Carrot pomace	5	28.49	50	0.570
	10	31.06	50	0.621
	15	34.23	50	0.685
	20	37.60	50	0.752
Potato peels	5	28.84	50	0.577
	10	30.89	50	0.618
	15	32.31	50	0.646
	20	34.95	50	0.699
Green pea peels	5	30.23	50	0.605
	10	35.43	50	0.709
	15	38.79	50	0.776
	20	41.01	50	0.820

Sensory evaluation of fiber substituted cakes at deferent levels.

The sensory evaluation of produced high fiber cake samples were evaluated. Twelve trained sensory panelists evaluated the products under investigation for their uniformity, size of cells, thickness, grain, moistness, tenderness, softness, crumb color, flavor and overall acceptability. The mean values were statistically analyzed using analysis of variance and least significant difference (LSD). In general, cake samples were significantly

different from control sample for all sensory properties. The results indicated that, the replacement of wheat flour caused a significant decrease in cake properties scores. The results in Table (8) showed that all high fiber substituted cake samples were significantly lower than control cake sample in all sensory characteristics. significant differences ($P>0.05$) in uniformity, size of cells, thickness, grain, moistness, tenderness, softness, crumb color, flavor and overall acceptability characteristics with control cake sample.

Table (8): Effect of addition dietary fibers sources on sensory attributes of produced cakes (mean±SD).

Dietary fiber sources	Substituted levels	Sensory attributes									
		Uniformity (10)	Size of cells (10)	Thickness (10)	Grain (20)	Moistness (10)	Tenderness (10)	Softness (10)	Crumb color (10)	Flavor (10)	Overall acceptability (100)
Control	0	9.65 ^a ±0.19	9.10 ^{ab} ±0.10	9.75 ^a ±0.47	19.60 ^a ±0.29	9.45 ^{ab} ±0.13	9.60 ^a ±0.11	9.70 ^a ±0.30	9.30 ^{ab} ±0.18	9.75 ^a ±0.16	97.65 ^{ab} ±1.95
Orange waste	5	9.60 ^a ±0.28	9.00 ^{ab} ±0.14	9.50 ^{ab} ±0.14	19.10 ^{ab} ±0.26	9.10 ^{ab} ±0.15	9.20 ^{ab} ±0.18	9.70 ^a ±0.20	9.40 ^{ab} ±0.16	9.30 ^{ab} ±0.18	97.20 ^{ab} ±1.55
	10	9.05 ^{ab} ±0.21	8.90 ^{ab} ±0.20	9.10 ^{ab} ±0.36	18.80 ^{ab} ±0.28	8.95 ^{ab} ±0.21	8.95 ^{ab} ±0.18	9.35 ^{ab} ±0.21	9.15 ^{ab} ±0.25	8.95 ^{ab} ±0.27	96.40 ^{ab} ±1.88
	15	8.75 ^{bc} ±0.09	8.75 ^{ab} ±0.24	8.85 ^b ±0.17	18.40 ^{ab} ±0.48	8.60 ^b ±0.27	8.70 ^b ±0.15	8.90 ^{ab} ±0.32	8.85 ^{ab} ±0.35	8.80 ^{ab} ±0.13	93.60 ^{bc} ±1.92
	20	7.80 ^c ±0.44	7.85 ^{bc} ±0.32	7.95 ^c ±0.21	17.60 ^{bc} ±0.27	7.80 ^{bc} ±0.22	7.70 ^c ±0.41	7.70 ^{bc} ±0.21	7.60 ^b ±0.26	7.80 ^{bc} ±0.21	88.90 ^{cd} ±2.02
Carrot pomace	5	9.50 ^{ab} ±0.41	9.30 ^a ±0.71	9.45 ^{ab} ±0.40	19.20 ^{ab} ±0.20	9.70 ^a ±0.84	9.40 ^{ab} ±0.32	9.45 ^a ±0.52	9.65 ^a ±0.42	9.65 ^{ab} ±0.33	97.95 ^a ±2.42
	10	9.05 ^{ab} ±0.37	9.00 ^{ab} ±0.47	9.10 ^{ab} ±0.38	18.80 ^{ab} ±0.39	9.15 ^{ab} ±0.72	9.00 ^{ab} ±0.36	9.05 ^{ab} ±0.79	9.05 ^{ab} ±0.62	8.75 ^{ab} ±0.31	93.35 ^{bc} ±2.27
	15	8.15 ^{bcg} ±0.28	8.20 ^b ±0.41	8.15 ^b ±0.34	18.00 ^b ±0.40	8.20 ^{bc} ±0.37	8.10 ^{bc} ±0.32	8.40 ^b ±0.32	8.55 ^b ±0.91	8.60 ^b ±0.34	90.30 ^c ±2.35
	20	7.10 ^{cd} ±0.38	7.60 ^{bc} ±0.65	7.45 ^c ±0.57	17.10 ^{bc} ±0.41	7.55 ^c ±0.32	7.40 ^{cd} ±0.39	7.50 ^{bc} ±0.67	7.45 ^c ±0.29	7.25 ^c ±0.47	86.35 ^d ±2.13
Potato peels	5	9.05 ^{ab} ±0.27	8.65 ^{ab} ±0.38	9.05 ^{ab} ±0.42	18.10 ^b ±0.22	9.00 ^{ab} ±0.51	9.10 ^{ab} ±0.48	8.75 ^{ab} ±0.37	8.40 ^b ±0.58	9.10 ^{ab} ±0.34	95.15 ^{ab} ±2.55
	10	8.50 ^{bc} ±0.24	8.00 ^{bc} ±0.38	8.40 ^b ±0.44	17.60 ^{bc} ±0.21	8.10 ^{bc} ±0.19	8.20 ^{bc} ±0.12	8.10 ^{bc} ±0.50	8.10 ^b ±0.10	8.30 ^{bc} ±0.64	88.90 ^{cd} ±2.02
	15	7.45 ^{cd} ±0.45	7.60 ^{bc} ±0.34	7.55 ^c ±0.29	16.40 ^c ±0.28	7.50 ^c ±0.31	7.40 ^{cd} ±0.54	7.20 ^c ±0.37	7.40 ^c ±0.38	7.20 ^c ±0.67	81.80 ^e ±2.46
	20	6.30 ^d ±0.63	6.20 ^d ±0.34	6.60 ^d ±0.43	15.70 ^c ±0.53	6.40 ^d ±0.53	6.80 ^d ±0.64	6.40 ^c ±0.52	6.60 ^c ±0.59	6.45 ^c ±0.56	79.50 ^e ±2.76
Green pea peels	5	9.10 ^{ab} ±0.24	9.00 ^{ab} ±0.48	9.70 ^a ±0.22	19.10 ^{ab} ±0.28	9.25 ^a ±0.32	9.00 ^{ab} ±0.41	9.25 ^{ab} ±0.35	9.10 ^{ab} ±0.58	9.15 ^{ab} ±0.29	94.10 ^b ±1.99
	10	8.25 ^{bc} ±0.55	8.40 ^{ab} ±0.62	8.10 ^b ±0.45	17.40 ^{bc} ±0.51	8.10 ^{bc} ±0.28	8.20 ^{bc} ±0.39	8.40 ^b ±0.17	8.50 ^b ±0.28	8.40 ^{bc} ±0.24	89.45 ^{cd} ±2.88
	15	7.00 ^d ±0.61	7.20 ^c ±0.33	7.60 ^c ±0.40	16.30 ^c ±0.63	7.30 ^c ±0.51	7.50 ^{cd} ±0.60	7.50 ^{bc} ±0.58	7.45 ^c ±0.51	7.45 ^c ±0.71	84.75 ^{de} ±2.85
	20	6.85 ^d ±0.68	7.00 ^{cd} ±0.61	7.10 ^d ±0.24	15.20 ^c ±0.24	6.90 ^{cd} ±0.32	6.95 ^{cd} ±0.34	7.00 ^c ±0.40	6.95 ^c ±0.39	7.05 ^c ±0.30	81.90 ^e ±2.46
L.S.D at $P<0.05$		0.79	0.91	0.84	1.43	0.86	0.89	1.02	1.07	1.14	3.72

^{a,b} There is no significant different ($P > 0.05$) between any two means, within the same attribute have the same letter

The results indicated that replacement of fiber with wheat flour substitutes caused a significant decrease in cake properties. However, there

were significant differences between cake samples containing the same type of fiber source at 5, 10, 15 and 20% replacement levels. The highest score in the same type of fiber source for all attributes were that achieved by cake samples with fiber source at 5% replacement level. On the other hand, the lowest scores in the same type of fiber source for all attributes were that achieved by cake samples with fiber source at 20% replacement level. For overall acceptability. Also for the all sensory attributes the cake samples at 0% substitution level (cake control) had the highest score, followed by 5% substitution level of all treatments for all attributes, then 10% substitution level for every type of fiber sources in all sensory characteristics. The cake samples at 20% substitution level had the lowest score. Samples prepared by using orange waste and carrot pomace had a slight higher score, but the lowest score were for samples prepared by potato and green pea peels. The above results are in agreement with those obtained by **Doweidar (2001)**; **El-Hadidi (2006)** and **Saeed (2010)** who reported that statistical analysis of sensory evaluation of the control cakes versus cakes containing some fibre powders showed insignificant differences.

Effect of addition dietary fiber sources on physical properties of produced cakes:

Data presented in Table (9) show that, the addition of dietary fiber sources was increased volume of cake under investigation, From this results indicated the important of adding dietary fiber sources on the volume of cake. Also, addition of dietary fiber sources was increased specific volume. The trend of increasing in the specific volume was go high after adding dietary fiber sources. These results are in agreement with **Saeed (2010)**.

Table (9): Effect of addition dietary fiber sources on physical properties of produced cake.

Dietary fiber sources	Substituted levels (%)	Weight (g)	Volume (cm ³)	Specific volume (cm ³ /g)
Control (Wheat flour 72%)	0	454.95	1084.93	2.385
Orange waste	5	455.68	1174.52	2.578
	10	456.91	1254.37	2.745
	15	458.49	1343.85	2.931
	20	461.12	1481.28	3.212
Carrot pomace	5	454.47	1127.58	2.481
	10	456.69	1245.91	2.728
	15	457.82	1289.07	2.816
	20	458.37	1357.19	2.961
Potato peels	5	457.88	1097.57	2.397
	10	461.45	1174.32	2.545
	15	464.58	1203.85	2.591
	20	469.37	1286.59	2.741
Green pea peels	5	452.18	1182.63	2.615
	10	453.64	1257.37	2.772
	15	452.87	1328.06	2.933
	20	453.93	1391.24	3.065

CONCLUSIONS

This study demonstrated the feasibility of using some food processing wastes to produce dietary fiber powders which may be used as a food ingredient. The results showed that the sources of dietary fiber had significant effects on the dietary fiber composition and technological properties. Moreover, the high effect on hydration properties which would affect the further application in real food system. Furthermore, carrot and orange fibers had high WHC and SWC values, which are good for food applications. Overall, the results suggested that orange waste, carrot pomace, potato peels and green pea peels could be used as a good raw material to produce dietary fiber powders. From industrial point of view, the fiber sources which are the residues from processing could be further processed to add value to the products in Egypt.

Future trends

Used new raw materials by-products, which containing a high amount of soluble and insoluble fractions of fibers and associated bioactive compounds. A good control of the parameters involved in this process (mainly in the washing and drying steps) could also be taken into account in order to minimize losses in the soluble dietary fiber components and in the biological activity of the associated compounds.

Acknowledgement

This research was partially funded by the Saudi Society for Food and Nutrition (SSFN). We thanks the Saudi Society for Food and Nutrition for this fund.

REFERENCES

- A.A.C.C. (2000). Approved methods of American Association of Cereal Chemists (10th ed.). published by American Association of Cereal Chemists, Ins. Saint Paul, Minnesota, USA.
- Abd El-Moniem, G.M. and Yassen, A.A.E. (1993). High dietary fiber cookies from several sources of bran or husk. Egypt. J. Food. Sci., 21: 157-170.
- A.O.A.C. (2000). Official Methods of Analysis (17th ed.). Arlington, VA: Association of Official Analytical Chemists, AOAC International.
- Camire, M.E. and Flint, S.I. (1991) Thermal processing effects on dietary fiber composition and hydration capacity in corn meal, oat meal, and potato peels. Cereal Chem. 68: 645–647.
- Camire, M.E.; Violette, D.; Dougherty, M.P. and McLaughlin, M.A. (1997) Potato peel dietary fiber composition: Effects of peeling and extrusion cooking processes. J Agric Food Chem., 45: 1404–1408.
- Carle, R.; Keller, P.; Schieber, A.; Rentschler, C.; Katzschner, T.; Rauch, D.; *et al.* (2001). Method for obtaining useful materials from the by-products of fruit and vegetable processing. Patent application, WO 01/78859 A1.
- Carme, G.M.; Susana, S.; Carmen, R. and Antoni, F. (2007). Effect of air during temperature on physicochemical properties of dietary and

- antioxidant capacity of orange (*Citrus aurantium* v. *Canoneta*) by-products. *Food chemistry*, 104: 1014-1024.
- Chantaro, P.; Devahastin, S. and Chiewchan, N. (2008). Production of antioxidant high dietary fiber powder from carrot peels. *LWT - Food Science and Technology* 41, 1987-1994.
- Chau, C. F., and Huang, Y. L. (2003). Comparison of the chemical composition and physicochemical properties of different fibers prepared from the peel of *Citrus sinensis* L. Cv. Liucheng. *Journal of Agricultural and Food Chemistry*, 51, 2615–2618.
- Doweidar, M. M. (2001). Chemical and physical studies on some natural resources used in improving bakery products. Ph. D. Thesis, Fac. of Agric, Cairo Univ., Egypt.
- El-Hadidi, S.T. (2006). Chemical and biological Evaluation of Cake products containing resistant starch and dietary fiber. Ph.D. Thesis, Fac. of Agric, Cairo Univ., Egypt.
- El-Refai, A.A.; El-Bastawesy, A. and Zakaria, M.M. (2006). Evaluation of some food processing wastes as sources of dietary fibers. *J. Agric. Sci. Mansoura Univ.* 31: 6505-6515.
- Esteller, M.S., Amaral, R.L. and Lannes, S.C. (2004). Effect of sugar and fat replacers on the texture of baked goods. *J. Texture Studies*, 35: 383–393.
- Femenia, A.; Lefebvre, C.; Thebaudin, Y.; Robertson, J. and Bourgeois, C. (1997). Physical and sensory properties of model foods supplemented with cauliflower fiber. *Journal of Food Science*, 62(4), 635–639.
- Fernandez-Gines, J. M.; Fernandez-Lopez, J.; Sayas-Barbera, E. and Perez-Alvarez, J. A. (2003). Effects of storage conditions on quality characteristics of bologna sausages made with citrus fiber. *Journal of Food Science*, 68(2), 710–715.
- Fernandez-Lopez, J.; Sendra-Nadal, E.; Navarro, C.; Sayas, E.; Viuda-Martos, M. and Pérez-Alvarez, J.A. (2009). Storage stability of a high dietary fibre powder from orange by-products. *International Journal of Food Science and Technology* 44, 748–756.
- Figuerola, F.; Hurtado, M.L.; Estévez, A.M.; Chiffelle, I. and Asenjo, F. (2005). Fibre concentrates from apple pomace and citrus peel as potential fibre sources for food enrichment. *Food Chemistry*, 91, 395–401.
- Gallaher, D. and Schneeman, B.O. (2001). Dietary fiber. In: B. Bowman and R. Russel (Eds.). *Present knowledge in nutrition* (8th ed., 805p). Washington, DC: ILSI.
- Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedures for Agriculture Research*. John Wiley and Sons Editor Inc. USA, 2 ed., Chapter 3, 129-184.
- Gomez, M.; Ronda, F.; Coballera, A.P.; Blanco, A.C. and Rosell, C.M. (2007). Functionality of different hydrocolloids on the quality and shelf life of yellow layer cakes. *Food Hydrocolloids*, 21(2), 167–173.
- Gorinstein, S.; Zachwieja, Z.; Folta, M.; Barton, H.; Piotrowicz, J.; Zember, M.; Weisz, M.; Trakhtenberg, S. and Martin-Belloso, O. (2001). Comparative content of dietary fiber, total phenolics, and minerals in

- persimmons and apples. *Journal of Agricultural and Food Chemistry*, 49, 952–957.
- Grigelmo-Miguel, N. and Martin-Belloso, O. (1999). Influence of fruit dietary fiber addition on physical and sensorial properties of strawberry jams. *Journal of Food Engineering*, 41, 13–21.
- Hassan, F.A.; Ismail, A.; Hamid, A.A.; Azlan, A. and Al-sheraji, S.H. (2011). Characterisation of fibre-rich powder and antioxidant capacity of *Mangifera pajang* K. fruit peels. *Food Chemistry*, 126, 283–288.
- Herbafood. (2002). Herbacel AQ Plus. Apple and citrus fibre. Available from www.herbafood.de/eaqplus.pdf. Date November 2002.
- Jaime, L.; Molla, E.; Fernandez, A.; Martin-Cabrejas, M.; Lopez Andreu, F. and Esteban, R. (2002). Structural carbohydrates differences and potential source of dietary fiber of onion (*Allium cepa* L.) tissues. *Journal of Agricultural and Food Chemistry*, 50, 122–128.
- Jimenez, A.; Rodriguez, R.; Fernandez-Caro, I.; Guillen, R.; Fernandez-Bolanos, J. and Heredia, A. (2000). Dietary fibre content of table olives processed under different European styles: Study of physicochemical characteristics. *Journal of the Science of Food and Agriculture*, 80, 1903–1908.
- Kethireddipalli, P.; Hung, Y.C.; Phillips, R. O. and McWatters, K. H. (2002). Evaluating the role of cell material and soluble protein in the functionality of cowpea (*Vigna unguiculata*) pastes. *Journal of Food Science*, 67(1), 53–59.
- Kuniak, L. and Marchessault, R. H. (1972). Study of crossing-linking reaction between epichlorhydrin and starch. *Starke*, 4, 110-116.
- Lamghari, R.; Sanchez, C.; El Boustani, E.; Maucourt, N. M.; Sauvaire, Y.; Mejean, L. and Villaume, C. (2000). Comparison of effects of prickly pear (*Opuntia ficus indica* sp.) fruits, arabic gum and citrus pectin on viscosity and in vitro digestibility of casein. *Journal of the Science of Food and Agriculture*, 80, 359–364.
- Lario, Y.; Sendra, E.; García-Pérez, J.; Fuentes, C.; Sayas-Barberá, E., Fernández-López, J. and Pérez-Alvarez, J.A. (2004). Preparation of high dietary fibre powder from lemon juice by-products. *Innovative Food Science and Emerging Technology* 5, 113–117.
- Larrauri, J. A. (1999). New approaches in the preparation of high dietary fibre powders from fruits by-products. *Trends in Food Science and Technology*, 10, 3–8.
- Larrauri, J.A.; Borroto, B. and Crespo, A.R. (1997). Water recycling in processing orange peel to a high dietary fibre powder. *Int. J. Food Sci. Tech.*, 32: 73–76.
- Lee, S. C.; Prosky, L. and DeVries, J. W. (1992). Determination of total, soluble and insoluble fiber foods. Enzymatic-Gravimetric method, Mes-Tris buffer. Collaborative study. *Journal of the Association of Official Analytical Chemists*, 75(3), 395–416.
- Mullin, W.J.; Wolunetz, M.S.; Emery, J.P. and Brooks, L. (1993). The effect of variety, growing location and storage on the dietary fiber content of potatoes. *J. Food Composition and Analysis*, 6: 316-323.

- Raghavendra, S.N.; Ramachandra-Swamy, S.R.; Rastogi, N.K., Raghavarao, K.S.M.S.; Kumar, S. and Tharanathan, R.N., (2006). Grinding characteristics and hydration properties of coconut residue: a source of dietary fibre. *Journal of Food Engineering*, 72, 281–286.
- Ralet, M.C.; Della Valle, G. and Thibault, J.F. (1993). Raw and extruded fibre from pea hulls. I. Composition and physico-chemical properties. *Carb Polymers*, 20: 17–23.
- Redgwell, R. J. and Fischer, M. (2005). Dietary fiber as a versatile food component: An industrial perspective. *Molecular Nutrition and Food Research*, 49, 521–535.
- Robertson, J. A.; Monredon, F. D.; Dysseleer, P.; Guillon, F.; Amado, R. and Thibault, T. F. (2000). Hydration properties of dietary fiber and resistant starch: a European collaborative study. *Lebensmittel Wissenschaft und Technologie*, 33, 72-79.
- Rodriguez, R.; Jimenez, A.; Guillen, R.; Heredia, A. and Fernandez-Bolanos, J. (1999). Postharvest changes in white asparagus during refrigerated storage. *Journal of Agricultural and Food Chemistry*, 47, 3551–3557.
- Rodriguez, R.; Jimenez, A.; Fernandez-Bolanos, J.; Guillen, R. and Heredia, A. (2006). Dietary fibre from vegetable products as source of functional ingredients. *Trends in Food Science and Technology* (17) 3–15.
- Saeed, M. A. (2010). Food processing for catering in spas. Ph.D. Thesis, Food Science Dept., Fac. of Agric., Moshtohor, Benha Univ., Egypt.
- Saleh, W.A.; Sharoba, A.M.; Bahlol, H.E. and El-Tanahy, H.H. (2012). Studies on Texture of some Foods. LAP LAMBERT Academic Publishing GmbH & Co. KG, Saarbrucken, Deutschland.
- Sanchez-Zapata, E.; Fernandez-Lopez, J.; Penaranda, M.; Fuentes-Zaragoza, E.; Sendra, E.; Sayas, E. and Perez-Alvarez, J.A., (2011). Technological properties of date paste obtained from date by-products and its effect on the quality of a cooked meat product. *Food Research International*, 44:2401–2407.
- Sanchez-Zapata, E.; Fuentes-Zaragoza, E.; Fernandez-Lopez, J.; Sendra, E.; Sayas, E.; Navarro, C. and Pérez-Alvarez, J.A. (2009). Preparation of dietary fibre powder from tiger nut (*Cyperus esculentus*) milk (“horchata”) by-products and its physicochemical properties. *Journal of Agricultural and Food Chemistry* 57, 7719–7725.
- Schieber, A.; Stintzing, F. C. and Carle, R. (2001). By-products of plant food processing as a source of functional compounds-recent developments. *Trends in Food Science and Technology*, 12, 401–413.
- Sharoba, A.M.; El-Mansy, H.A. and Senge, B. (2012). Rheological and Mechanical Properties of Some Selected Foods. LAP LAMBERT Academic Publishing GmbH & Co. KG, Saarbrucken, Deutschland.
- Sosulski, F.W. and Wu, K.K. (1988). High-fiber breads containing field pea hulls, wheat, corn and wild oat brans. *Cereal Chem.*, 65: 186–191.
- Sowmya, M.S.; Jeyarani, T.; Jyotsna, R. and Indrani, D. (2009):Effect of replacement of fat with sesame oil and additives on rheological, microstructural, quality characteristics and fatty acid profile of cake. *Food Hydrocolloids*, 23: 1827–1836.

- Sudha, M.L.; Baskaran, V. and Leelavathi, K., (2007). Apple pomace as a source of dietary fibre and polyphenols and its effect on the rheological characteristics and cake making. *Food Chemistry*, 104, 686–692.
- Toma, R.B.; Orr, P.H.; D'Appolonia, B.; Dintzis, F.R. and Tabekhia, M.M. (1979). Physical and chemical properties of potato peel as a source of dietary fiber in bread. *Journal of Food Science*, 44, 1407–1417.
- Tudorica, C. M.; Kuri, V. and Breenan, C. S. (2002). Nutritional and physicochemical characteristics of dietary fiber enriched pasta. *Journal of Agricultural and Food Chemistry*, 50, 347–356.
- Vazquez-Ovando, A.; Rosado-Rubio, G.; Chel-Guerrero, L. and Betancur-Ancona, D. (2009). Physicochemical properties of a fibrous fraction from chia (*Salvia hispanica* L.). *Food Science and Technology*, 42, 168–173.
- Wilkins, M.R.; Suryawati, L.; Maness, N.O. and Chrz, D. (2007). Ethanol production by *Saccharomyces cerevisiae* and *Kluyveromyces marxianus* in the presence of orange-peel oil. *World J. Microbiol. Biotechnol.*, 23 (8), 1161–1168.

استخدام مخلفات تصنيع بعض الخضراوات والفاكهة كمصدر للألياف الغذائية في صناعة الكيك

أشرف مهدي عبدالحميد شروبه* ، منى عبدالرحيم فراج** و
أميرة محمد عبدالخالق عبدالسلام**
* قسم علوم الأغذية – كلية الزراعة بمشتهر – جامعة بنها
** معهد بحوث تكنولوجيا الأغذية – مركز البحوث الزراعية

تم في هذه الدراسة تجهيز بعض مخلفات تصنيع الخضراوات والفاكهة لإستخدامها كمصدر للألياف الغذائية في التصنيع الغذائي. تم تجهيز مخلفات كل من البرتقال والجزر والبطاطس والبسلة الخضراء وتم دراسة الخواص الوظيفية للألياف والتركيب الكيماوي لها. وتم إستخدامها بعد ذلك في إنتاج كيك منخفض السعرات الحرارية حيث تم دراسة تأثير أستبدال الدقيق إستخلاص ٢٢% المستخدم في صناعة الكيك بمخلفات تصنيع هذه الخضراوات والفاكهة بعد تجفيفها كمصدر للألياف الغذائية بنسبة أستبدال ٥، ١٠، ١٥ و ٢٠%. وذلك لخفض السعرات الحرارية ورفع نسبة الألياف للكيك لكي يلانم احتياجات مرضى السمنة ومرضى السكر وراغبي انقاص الوزن. وقد تمت دراسة تأثير هذه الاستبدالات على كل من الخواص الفيزيائية والتركيب الكيماوي و الصفات الحسية للكيك وذلك بجانب دراسة الصفات الريولوجية لخلطات الدقيق والألياف وقياس ثوابت القوام للكيك الناتج من هذه الخلطات. وقد أظهرت النتائج المتحصل عليها أن الخواص الريولوجية لدقيق القمح تأثرت وزادت نسبة الإمتصاص للماء وكذلك زيادة زمن العجن نتيجة الاستبدال بمصادر الألياف مقارنة بالعينة القياسية، وقد زادت فترة الثبات للعينات المحتوية على مخلف الجزر ومخلفات تصنيع البرتقال وقشور البطاطس وقشور البسلة الخضراء بينما أنخفضت قليلا درجة الضعف للعينات المحتوية على مصادر الألياف. وأدت إضافة مصادر الألياف إلي انخفاض في الحجم والحجم النوعي للكيك مقارنة بالعينة القياسية. حقق مخلف الجزر أفضل النتائج الحسية وجاءت مخلفات تصنيع البرتقال في المرتبة الثانية ثم تلتها قشور البطاطس وقشور البسلة الخضراء لجميع نسب الإستبدال المختلفة، وأظهرت نتائج التقييم الحسي للكيك أن جميع نسب الإستبدال مقبولة حسيا ماعدا بعض الصفات الحسية كانت غير مرضية عند نسب استبدال ٢٠% لجميع مصادر الألياف المستخدمة، وكانت أفضل النتائج لعينات الكيك المحتوية على مخلف الجزر ومخلف البرتقال بنسب استبدال ٥ و ١٠% على التوالي.

قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة
كلية الزراعة – جامعة بنها

أ.د / احمد عبد العزيز الرفاعي
أ.د / حمدي عبد اللطيف المنسي

