

## **EFFECT OF HUMAN NUTRITION BY MEALS CONTAINING GHEE, OLIVE OIL, AND A BLEND OF THEM ON SERUM LIPIDS.**

**Abdeen, E.M. I. <sup>1</sup>; A. S. Abd - El-Star <sup>2</sup>; A. A. El-Cerss<sup>3</sup>; A. T. Gaber<sup>4</sup>; M. Abd-El- Fatah<sup>2</sup> and Awatif I. Ismail<sup>2</sup>**

**1- Pasta and Bread Res. Dept. (FTRI), (ARC), Giza, Egypt.**

**2- Oil and Fat Res. Dept. (FTRI), (ARC), Giza, Egypt.**

**3- Physical Education, Tanta Univ. Egypt.**

**4- Dairy Products Res. Dept. (FTRI), (ARC), Giza, Egypt.**

### **ABSTRACT**

This research was conducted for the lack of the quantity of corn sufficient to obtain the pure corn oil which are found in a large quantity in the market. The reason is that embryo weight represents 4% of the weight of corn grain and the proportion of corn oil in embryo 25-27 % of its weight. So tons of maize needed to get 10 kilograms of oil, if the oil extraction rate 100% and this is impossible. Also some investigators, showed that corn oil and other industrial seed oils promote strongly prostate cancer cell growth and increase mortality.

As well as the desire of consumers to ghee nutrition because of their taste , distinctive flavor and beneficial effects on brain cells and afraid at the same time a high proportion of cholesterol in the blood.

Where experiments were carried out on nutrition human where meals are contained ghee with olive oil each separately and mix them as a source of fats (25: 29 % of total energy). The analysis of the fatty acids present in olive oil and buffalo ghee, showed that olive oil was free of cholesterol's and short chain fatty acids (4:15 C). While, ghee contained 180 mg/dl and short chain fatty acids.

The design of experiments has been developed on the basis of three variables, each variable at three levels as following:

- 1 – The practice of physical activity, at three levels (do not engage in any activity – without regular exercise – exercise regularly). By a proposed training program for the development of the element of balance and stability during movement in the compulsory kata karate. The program includes a number of physical techniques and exercises to build muscles of the upper limb (forearm) for fist and lower limb (thigh) kicked. And skills necessary for a group that practiced karate on a regular basis. And subjected to a diet program, Faculty of Physical Education - University of Kafr-El-Sheikh.
- 2 – The age group, and there levels are (less than 20 years old – from 20:30 years old – more than 30 years old).
- 3 – The obesity, and there levels are (thin – average – fat).

The blood sample was drawn from all individuals participating in the experiment and assess triglyceride (TG) , low lipoprotein (LDL-c) , high (HDL-c) density and total cholesterol (TC) and the exclusion of non-matching samples (Referencerange).

Diet Nutrition (based on 30% of energy from fat) conducted for three consecutive weeks and then a blood sample was withdrawn and another week after feeding. And calculate the averages for each of the TG, LDL-c, HDL-c, and TC.

The results found that feeding diets containing ghee raise the proportion of about 7% cholesterol and reduce triglycerides .The practice of sports activities and age group less than 20 years old and medium weight less influence from the other . Feeding diets containing olive oil reduces cholesterol including high density lipoprotein

cholesterol and raise the proportion of TG, which may be also a risk to public health . The practice of physical activity reduces triglycerides and reduce cholesterol, especially low-density.

Diets containing a mixture of ghee and olive oil did not cause change , both in the level of cholesterol or triglycerides for individuals who engage in sports activities. On this basis, feeding is a mixture of ghee and olive oil feeding healthy, safe, to avoid the diseases of blood, heart and arteries, especially when the practice of physical activity for the building blocks of muscles and overcoming. The good of vegetable oils in general with the statement ratios between omega-6 and omega-3 and modified by mixing vegetable oil with each other to become the ratio between omega-6: omega-3 is 4: 1 to amend any mixing ratios to avoid the risk of cancer.

## **INTRODUCTION**

This research was conducted for the lack of the quantity of corn sufficient to obtain pure corn oil , which are found in a large quantity in the market. Our previous studies on 8 corn varieties which cultivated in Egypt namely single hyper named (Giza129and Bashaier13 as white varieties – Pioneer and Giza 155 as yellow corn varieties). Three hyper named (Wattania and Nevertity as white –Giza 351and 352 as yellow corn), showed that corn embryo weight represents 4% of the weight of corn grain and the proportion of corn oil in embryo represents 25-27 % of its weight. So , tons of maize needed to get 10 kilograms of oil, if the oil extraction rate 100% and this is impossible.

Also some investigators, showed that corn oil and other industrial seed oils promote strongly prostate cancer cell growth and increase mortality.

In fact, there have even been several experiments showing that butter and other animal fats promote cancer growth to a lesser degree than margarine and omega-6-rich seed oil. A high intake of omega-6 fatty acids may increase the likelihood that postmenopausal women will develop breast cancer (Emily, *et al.*, 2008) .Similar effect was observed on prostate cancer (Yong, *et al.*, 2007). Other analysis suggested an inverse association between total polyunsaturated fatty acids and breast cancer risk (Valeria, *et al.*, 2001).Ghee is widely used in food habits of Arabic zone. Ghee supports the functioning of the brain. Excessive mental work can aggravate Prana Vata, which is responsible for creative thinking and reasoning. PranaVata, when out of balance, will dry out Tarpaka Kapha, the dosha that lubricates the brain. Cooking of ghee regularly considered a rasayana is its medhya quality supports the moisturizing effect of Tarpaka Kapha (Ayurveda, 2006). Ghee also, enhances the quality and quantity of Ojas, the master coordinator of mind are necessary for the development of our full human potential or enlightenment. The proper amount of ghee also enhances Agni (Digestive fire). But too much ghee has think of a ghee lamp: in order to burn it you need to keep adding ghee. Ghee, which is rich in saturated fatty acids it produced from butter of cow or buffalo milk fat. Because ghee has such a high smoke point (252 °c) it is a very useful oil for cooking. The smoke point determines when oil actually starts to burn and generate oxidation and the potential free radicals. As it has a very low oxidation rate ghee stays fresh even UN - refrigerated for a long time. Another benefit of using ghee is that the heating

procedure removes the lactose content making it tolerable to those sensitive to lactose (Ayurveda, 2006).

Valeria, *et al.* (2001) reported that the type of fatty substances is major factor of blood cholesterol level (solid fats raise this level, while oils lower it).

This research was conducted for the purpose of the lack of quantities corn dominating for pure corn oil.

On other hand dairy products make an appreciable contribution to saturated fat and cholesterol intake. Consumption these products may be correlated with high blood cholesterol level, ( Rossouw, *et al.*, 1981).

High blood cholesterol is the major risk factor for cerebrovascular disease (NCEP.1993).

Through a period of time many research workers showed a direct relation between cholesterol and the occurrence of coronary heart disease, (John , *et al.* 1990 and Rawashdeh, 2002). There are two types of lipoproteins (low and high density lipoproteins) low density lipoproteins are responsible for segmentation cholesterol in the artery walls, and its concentration is associated with a higher risk of heart attack. High density lipoprotein concentration may play a beneficial role (Koo, *et al.* 1985). Transport cholesterol from cells to the liver for reprocessing or bile acids formation (Hui, 1992).

On contrary recent studies show serious health problems when feeding on some vegetable oils such as the famous corn oil.

Corn agronomists have developed high oil varieties; however, these varieties tend to show lower field yields, so they are not universally accepted by growers, Dupont, *et al.* (1990).

Refined corn oil constituents are 99% triglyceride, with proportions of approximately 55% polyunsaturated fatty acid, 30% monounsaturated fatty acid, and 15% saturated fatty acid. Of the saturated fatty acids, 80% are palmitic acid (lipid number of C 16:0), 14 % stearic acid (C 18:0) and 3% arachidic acid (C 20:0). Over 99% of the monounsaturated fatty acids are oleic acid (C18:1c). 98% of the polyunsaturated fatty acids are the omega-6 linoleic acid (C18:2 n-6 c, c) with the 2% remainder being the omega-3 alpha-linolenic acid (C 18:3 n-3 c, c, c) (USDA, 2007).

Some medical researches suggest that excessive levels of omega- 6 fatty acids, relative to omega -3 fatty acids, may increase the probability of a number of diseases and depression .(lands, 2005 , Hibbeln ,*et al.*, 2006, and Okuyama , *et al.*, 2007 ) . Modern Western diets typically have ratios of omega -6 to omega-3 in excess of 10 to 1 , some as high as 30 to 1, partly due to corn oil which has an omega-6 to omega -3 ratio of 49 : 1 . The optimal ratio is thought to be 4:1 or lower (Daley, *et al.*, 2004 and Simopoulos 2002).

Objective of the research: is to find the perfect way to feed the plant oils and fat without a negative impact effect on public health. Do not rely on certain class oil, but after studying its components and modify the components by mixing other oils to avoid negative impacts on health.

Down on the willingness of consumers to nutrition ghee because of its taste and flavor is good and beneficial effects on brain cells. So, the study was conducted to determine the means of proper nutrition ghee. Also, this

study was interested with effects of buffalo ghee and olive oil upon Egyptian blood lipid profile. So for none of the studies reported in Egypt to assess the effect of ghee on serum lipid profile of human.

## MATERIALS AND METHODS

The experimental design was studied on human after announcing their relations according to sports activity, age's, and obesity.

Three independent variables at three levels as the reported design by Abdeen, and Agag (1999) as follows:

- 1-Sport activity: (Avoid for physical activity, Irregular, and Viol regularly).by impact of a proposed training program for the development of the element of balance and stability during movement in the compulsory kata karate. The program includes a number of physical techniques and exercises to build muscles of the upper limb (forearm) for fist and lower limb (thigh) kicked, a skills necessary for a group that practiced karate on a regular basis, and subjected to a diet program, Faculty of Physical Education - University of Kafr- El-Sheikh.
- 2- Age: (less than 20, 20– 30 and more than 30).
- 3- Obesity: (Skinny, Medium, and Fleshy).

**Table (1): The three variables at three levels variation, which experimental designed for human nutrition.**

Variables	Variation			
	-1	0	+1	Cod
Sport activity	Avoid for physical activity	Irregular	Viol regularly	A
Age	Less than 20 years	20- 30 years	More than 30 years	B
Obesity	Skinny	Medium	Fleshy	C

**Table (2): The out line of the experimental design.**

Ex. No.	Test condition							Variables							
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
A	-1	+1	-1	+1	-1	+1	-1	+1	0	0	0	0	0	0	0
B	-1	-1	+1	+1	0	0	0	0	-1	+1	-1	+1	0	0	0
C	0	0	0	0	-1	-1	+1	+1	-1	-1	+1	+1	0	0	0

### Diet :

Main source of carbohydrate:

Macaroni (two times a week)

Rice (three times a week )

Bread (three times a day )

### Main source of protein:

Chicken, eggs, milk and its products, meat, bean and lentils.

### Main source of fat:

Olive oil as row was used to add to bean and dairy products .

The only variable was the olive oil and buffalo Ghee .

Vegetable and fruits were eaten twice a week..

Buffalo ghee was used in cooking and dipping .

Buffalo ghee and olive oil were purchased from department of animal farm Sakh and local supermarket, respectively.

Fasting venous blood samples (10 ml) were collected at the entry and the end of period.

**The 45 subjects ate:** Their food with buffalo ghee or olive oil or mixed as the sole source of fat type for 3 wk., as description of diet composition Table (3).

**Table (3): Food consumption of subjects.**

Dietary fat type	Variables			Energy Calorie	Carbohydrate		Fat		Protein	
	A	B	C		g	%	g	%	g	%
Olive oil	-1	-1	-1	1800	298	66.7	50	25	40	8.8
	0	0	0	1980	320	64.6	60	27.2	45	9.1
	+1	+1	+1	2026	350	69.1	62	27.9	45	8.8
Buffalo ghee	-1	-1	-1	1800	290	64.4	49	24.5	40	8.8
	0	0	0	1980	320	64.6	60	27.2	45	9.1
	+1	+1	+1	2026	330	65.1	62	27.5	45	8.8
Blend	-1	-1	-1	1800	310	68.8	50	25.0	40	8.8
	0	0	0	1980	320	64.6	60	27.2	45	9.0
	+1	+1	+1	2026	330	65.1	62	27.2	45	8.8

Table (3) for a description of diet composition. The basic diet was formulated to provide 28% of food energy as fat. The actual range varied from 25 to 27.9 energy % because subjects were varied.

The subjects were encouraged to maintain a constant body weighted and were weighted daily.

All food was provided for the subjects as three meals per day and was prepared in a local bakery. Subjects used check lists to keep daily records of all food eaten, including any occasional extramural food consumption. A staff member regularly verified with the subjects that these records were complete and accurate. In this way , the overall composition of the diet stayed constant, only the type of lipid used in food preparation was changed. Food intake study was noted that olive oil consumed by volunteers more than other oil or fat. It was used at least twice a day, consistently for breakfast and lunch, especially with chickpea and thyme and many times used for cooking.

Ghee used as delicious oil for cooking or stirs into rice and vegetables for a nourishing flavor some taste.

However, the period 3wk was chosen as it had previously been shown that plasma lipid levels stabilized within 2:3 wk after initiating a change in dietary fat (Bonamon and Grundy, 1988). Fasting venous blood samples (10 ml) were collected at the entry and the end of period for analysis.

**Chemical analysis :**

**Preparation of the fatty acid methyl esters .**

According to the procedure of Morrison and Smith (1962) fatty acids composition of buffalo ghee and olive oil were saponified and the free fatty acids were methylated .

The methyl esters of olive oil and buffalo ghee was prepared using benzene, methanol and concentrated sulfuric acid (10 :86 :4) and methylation was carried out for one hour at 80 -90 ° C according to Stahl (1967)

**Identification of fatty acids methyl esters:**

Gas- liquid chromatography apparatus (Apye unicom model 4550) was used. The conditions were identical to those reported by Ishmael (1989). Peak areas were measured using spectrophysic integrator .

Total cholesterol of ghee and olive oil were determined according to the method of Plummer (1978).

Serum total cholesterol was determined using the enzymatic method at EI – Resala Lab. Kafr El-Shiekh.

Triglycerides (TG) were determined by the enzymatic technique from EI-Resala Lab. HDL-c was analyzed by the precipitation technique using magnesium chloride and phosphotungstic acid. LDL-c was calculated using the formula of Friedewald *et al.* (1972).

## **RESULTS AND DISCUSSION**

Saturated fatty acids (SFA), monounsaturated fatty acids (MUSFA), polyunsaturated fatty acids (PUSFA) and total cholesterol (TC).

The fatty acid profiles and total cholesterol content of olive oil and buffalo ghee in experimental food consumption are presented in Table (4). The results showed that olive oil was free of cholesterol and short chains fatty acids (4:15C) .While, buffalo ghee contained 180mg /dl and short chins fatty acids.

Ghee was rich in easy- to- digest short chain fatty acids. It also contained 4.5 % linoleic acid which had anti-oxidant properties . So, it is healthy cooking oil due to its content of essential fatty acids which are an essential part of our diets.

Ghee has a high smoke point which makes it great for cooking. It means, that it does not produce damaging free radicals .Elson (1992) reported that besides that ghee contained appreciable amounts of short and medium chain fatty acids (FAS), these FAS may be an activator for hepatic hydroxyl-methyl-glutrate-coA reductase, the rate limiting enzyme in cholesterol biosynthesis.

Olive oil contained the higher MUSFA and PUSFA (67.2% -12, 9 %), the lower SFA (19. 8%) than buffalo ghee. The ratio of PUSFA / SFA(P:s) and MUSFA +PUSFA (m + p) /s have been used by nutritionists to interpret the effect of dietary oil and fat on the level of blood cholesterol (Hodeson *et al.*, 2001). The results also indicated that olive oil was higher in this parameter than ghee . As seen in Table (5) The proximate analysis of the blood serum lipid concentrations of participants (TG,HDL-c, LDL-c and total cholesterol -TC) before starting of 3 wk feeding are shown in this Table . The Table shows the health status of individuals based on experience after excluding the results according to the reference rang.

**Table (4): Fatty acid profile of buffalo ghee and olive oil used in the experimental.**

Fatty acid	Buffalo ghee	Olive oil
4,6,8:0	6.6	----
10:0	6.0	----
12:0	4.2	----
14:0	8.3	----
14:1	1.3	----
14:2	0.6	----
15:0	0.3	----
16:0	24.4	14.6
16:1	2.8	0.7
16:2	0.3	0.2
17:0	1.1	----
18:0	10.3	5.2
18:1	28.2	66.5
18:2	4.5	12.0
18:3	0.8	0.7
20:0	0.3	----
SFA% (S)	61.5	19.8
MUSFA% (M)	32.3	67.2
PUSFA% (P)	6.2	12.9
P:S	0.10	0.65
M + p / s	0.62	4.04
TC (mg / dl)	180.0	----

**Table (5): Shows the results of the analysis for each of the blood TG, HDL-c, LDL-c and total cholesterol (TC) before beginning of the experimental .**

Exp. No:	TG mg/dl			HDL-c mg/dl			LDL-c mg/dl			TC mg/dl		
	35-165			> 46.32			< 135.1			150-200		
References rang	O.O	Ghee	blend	O.O	Ghee	blend	O.O	Ghee	blend	O.O	Ghee	blend
1	99	108	103	50	48	52	110	130	130	160	178	182
2	88	68	106	62	49	66	111	120	104	173	169	170
3	73	69	55	56	50	73	109	123	90	165	173	163
4	98	69	73	74	49	62	106	122	98	180	171	160
5	80	99	82	52	47	55	142	129	105	194	176	160
6	82	70	91	63	48	47	100	130	128	163	178	175
7	76	78	72	49	50	56	119	125	112	168	175	168
8	88	79	76	58	53	53	125	122	120	183	175	173
9	50	98	48	62	55	64	109	109	118	171	164	182
10	68	103	106	74	67	72	90	107	115	164	174	187
11	94	100	103	61	75	81	91	120	99	152	195	190
12	55	99	93	73	80	83	85	88	96	158	168	181
13	73	77	86	82	70	71	97	105	76	179	175	173
14	82	102	95	81	68	66	100	115	75	181	183	175
15	93	121	100	76	60	64	117	112	73	193	172	190

Serum lipid response of the participants at starting and final of the 3wk feeding of ghee are shown in Table (6).

Ghee raised serum TC level a bout 7.3% in average and LDL-c level (+ 7.2%) and caused an increase in HDL-c level about 5.8% in average above the initial levels. Among the variables ( A, B, and D) Exp. No. 1 ,5,7,and 12 caused the highest raise in serum TC levels . These subjective as shown in Table (1 and 2 ) avoid for physical activity , more than 30 years , and fleshy .

**Table (6): Serum lipid responses (mg/dl) and % from the base line of variables after 3 wk. fed dietary ghee.**

Exp. No.	TG		HDL-c		LDL-c		TC	
	mg/dl	%	mg/dl	%	mg/dl	%	mg/dl	%
1	-3	-2.7	+3	+6.2	+15	+11.5	+18	+8.5
2	-1	-1.4	+4	+8.1	+5	+4.1	+9	+5.3
3	-3	-4.3	+3	+6.0	+12	+9.7	+15	+8.6
4	-2	-2.8	+4	+8.1	+6	+4.9	+10	+5.8
5	-3	-3.0	+2	+4.2	+17	13.1+	+19	+10.7
6	-1	-1.4	+4	+8.3	+4	+3.1	+8	+4.4
7	-3	-3.8	+2	+4.0	+16	+12.8	+18	+10.2
8	-2	-2.5	+3	+5.6	+9	+7.3	+12	+6.8
9	-1	-1.0	+4	+7.2	+4	+3.6	+8	+4.8
10	-1	-.97	+4	+5.9	+5	+4.6	+9	5.1+
11	-2	-2.0	+2	+2.6	+12	+10.0	+14	+7.1
12	-3	-3.0	+3	+3.7	+13	+6.9	+16	+9.5
13	-2	-2.5	+4	+5.7	+6	+5.7	+10	+5.7
14	-2	-1.9	+4	+5.8	+7	+6.0	+11	+6.0
15	-2	-1.6	+4	+6.6	+6	+5.3	+10	+5.8

On the contrary Exp. No. (2,4,6,9,10,13,14, and 15 ) showed the lowest raise in serum TC levels . These subjects as shown in Table (2) are viol regularly sports, young , and skinny or medium . In general , serum cholesterol levels tended to be low reference value (150 :200 mg/dl) in all experimental .

Also, the results showed that reduction TG in all Exp. Subject .

If all standards of nutrition on ghee raise the proportion of total cholesterol and reduce the rate ruling on triglyceride. It was noted the increase in total cholesterol as a result of feeding ghee to the extent permitted and the increase in HDL-c in groups that viol regularly sports and small-and medium-Age Forums . Many nutrients other than dietary fat influence cholesterol concentrations such as low cholesterol concentration, dietary fiber, complex carbohydrate intake and protein source (Cohen *et al.*, 1994).

As expected, it was noted increase of the proportion of total cholesterol in general as a result of feeding ghee. However, high cholesterol for each of the sports practice, young and slim due to the increase in HDL-c. Returns while increasing the proportion of total cholesterol for each of the Sport and shall not perform advanced in age and fat due to an increase in LDL-c . Elson

(1992) reported that besides that ghee contained an appreciable amount of short and medium chain fatty acids (FAS), these FAS may be work as an activator for hepatic hydroxyl-methyl-glutrate-coA reductase, the rate limiting enzyme in cholesterol biosynthesis. Short chain FAS with two other FAS namely, lauric and myristic acids were thought as hypercholesterolenic agents. Ide *et al* . (1979) completed that and may be cause high rate of cholesterol absorption. Hajji *et al.*(1998) reported that myristic acid appeared to be the most potent cholesterol rising in human. Also, the cholesterol content of ghee may be increased blood cholesterol concentration (Gurr, 1989).

The results showed that a great reduction in the concentrating of serum total cholesterol, HDL-c and LDL-c and these reduction may be due to high MUSFA (67.2) oleic acid content in olive oil .

Polyphenolic compounds found in olive oil may be contributed to the lower incidence of coronary heart disease. Recent findings demonstrate that olive oil phenolics inhibit oxidation of LDL-c (Visioli and Galli, 1999) .

Garg, (1994) reported that olive oil for routine frying or cooking practices, that is it did not produce toxic aldehyde.

**Table (7): Serum lipid responses (mg/dl) and % from the base line of variables after 3 wk. fed dietary olive oil.**

Exp .No.	TG		HDL-c		LDL-c		TC	
	mg/dl%		mg/dl%		mg/dl%		mg/dl%	
1	+5	+4.6	-3	-2.3	-6	12.5-	-9	5.0-
2	+3	+4.4	1-	0.8-	3-	6.1-	4-	-3.2
3	+4	+5.9	3-	2.4-	6-	12.2-	9-	5.2-
4	+3	+4.3	3-	2.4-	5-	10.2-	8-	4.6-
5	+5	+5.0	3-	2.3-	5-	10.6-	9-	5.1-
6	+3	+4.2	1-	0.7-	3-	6.2-	-4	2.2-
7	+5	+6.4	3-	2.4-	6-	12-	-9	5.1-
8	+4	+5.0	2-	1.6-	5-	9.4-	7-	4.0-
9	+3	+3.0	2-	1.8-	4-	7.2-	6-	3.6-
10	+3	+2.9	1-	0.9-	3-	4.4-	4-	2.2-
11	+4	+4.0	2-	1.6-	3-	4.0-	5-	2.5-
12	+5	+5.0	2-	2.2-	7-	8.7-	9-	5.3-
13	+3	+3.8	1-	0.9-	3-	4.2-	4-	2.2-
14	+3	+2.7	1-	0.8	3-	4.4-	4-	2.1-
15	+3	+2.4	-1	0.8	3-	5.0-	4-	2.3-

Olive oil consumption lowered levels of serum total cholesterol (-3.64 %), LDL-c (-7.8%) and caused slight reduction in HDL-c (-1.9%) on average, under the initial level. On the contrary , olive oil raised TG (+3.7%) on average. Among the variables (A, B, and C), Exp. No. 6, 10, 11, 13, 14 and 15 serum TC level was decreased. These subjects as shown in Table (1-2) are viol regularly sports, young, and skinny or medium. Whereas, Exp. No. 1, 3, 4, 5,7and8 showed the highest level of serum TC. The results showed that olive oil caused reduction in the concentration of serum TC, LDL-C and HDL-C and this reduction may be due to high oleic acid content 6.6% in olive oil. Also polyphenolic compounds found in olive oil may be contributed to the

lower incidence of coronary heart disease in this area, (Rawashdeh, 2002). Visioli and Galli (1999) demonstrated that olive oil phenolics inhibit oxidation of low-density lipoproteins. Garg,( 1994 )reported that olive oil can be advised as an alternative to high –carbohydrate diets in diabetic and carbohydrate-sensitive patients. Also Grooviest, (1998) reported that olive oil did not produce toxic aldehydes during routine frying or cooking practices. Generally, olive oil feeding decrease TC, HDL-c, LDL-c and increase TG.

The presence of a high level of this fat (TG) especially with the presence of a low level of high density lipoprotein HDL-c in the blood that looks like a tremor of injury of heart disease, although it is not the same force, such as high cholesterol level, (Lands, 2005).

**Table (8): Serum lipid responses (mg/dl) and % from the base line of variables after fed dietary blend of ghee and olive oil.**

Exp. No.	TG		HDL-c		LDL-c		TC	
	mg/dl	%	mg/dl	%	mg/dl	%	mg/dl	%
1	+2	+1.8	+1	+0.76	+2	+4.1	+3	+1.7
2	-	-	-	-	-	-	-	0
3	+3	+4.3	+2	+1.6	+2	+4.0	+4	2.3+
4	-	-	-	-	-	-	-	0
5	+2	+2.0	+1	+0.77	+1	+2	+2	1.1+
6	-	-	-	-	-	-	-	0
7	+2	+2.5	+2	+1.6	+1	+2.0	+3	+1.7
8	-	-	-	-	-	-	-	0
9	+1	+1.0	+2	+1.8	+1	+1.8	+3	+1.8
10	+1	+9.0	+2	+1.8	+1	+1.4	+3	+1.7
11	+1	+1.0	+1	+0.8	+1	+1.3	+2	+1.0
12	+1	+1.0	+1	+1.1	+1	+1.2	+2	+1.2
13	+1	+1.2	-	-	-	-	-	0
14	+1	+0.9	-	-	-	-	-	0
15	+1	+0.8	-	-	-	-	-	0

It was shown that no change in serum TG, HDL-c, LDL-c and TC in EXP.NO. (2, 4, 6, and8) when blended ghee with olive oil was used . Also, EXP. NO. (13, 14, and15). A slight increase was shown in this parameter in the residual EXP.

Some studies showed that diets high in PUSFA can lower serum HDL-c concentration (Hajji *et al.*, 1998 and Hodson, *et al.*, 2001). This means that blend olive oil (PUSFA and MUSFA) with ghee (SFA) resulted a slight increase in serum parameter. The human body is 80% of the requirements of the cholesterol in the liver and the remaining 20% has to be dealt with sources of food for foreign affairs. (Garg, 1994).

The distained data suggest that this may not be absolutely necessary avoid ghee for those who are suffering heart diseases.

Finally: Olive oil which is high oleic acid, with ghee may be a good alternative of dietary fat for reducing blood cholesterol level.

## REFERENCES

- Abdeen, E.M.I, and Agag, M.E.M., ( 2002) . New products from rice application for 100% good quality rice bread without yeast. Arab Republic of Egypt ,Ministry of State for Scientific Research, Academy of Scientific Research and Technology, Technology Development and Scientific Services Sector, Patent Gazette (2002). No: 595.
- Ayurveda, A.I. (2006). The virtues of ghee. *Physical Nutrition*, 26 (3):1-3.
- Bonamon, A. and Grundy, S.M. (1988). Effect of dietary stearic acid on plasma cholesterol and lipoprotein levels. *N. Engl. J. Med.*, 318:1244-8.
- Cohen, B.I., Ayyad, N., Mikami, T., Mikami, Y. and Mosbacch, E.H. (1994). Effects of dietary fat and fatty acids on sterol balance in hamsters. *Lipids*, 29:503.
- Daley, C. A., Abbott, A., Doyle, P., Nader, G.and Larson, S. (2004). A literature review of the value- added nutrients found in grass-fed beef products. California State University, Chico (College of Agriculture).  
<http://www.csuchico.edu/agar/grassfedbeef/healthbenefits/insex.html:Retrieved> 2008-03-23.
- Dupont, J., White, P.J., Carpenter, M.P. Schaefer, E.J., Meydani, S.N., Elson, Woods, M., and Gorbach, S.L. (1990). Food uses and health effects of corn oil. *J .Am. Coll. Nutr.* 9 (5):438-470. PMID 2258533.
- Elson. E. (1992). Tropical oils nutritional and scientific issues. *Critical Reviews. Food Sic. Nutr.* 31:79-102.
- Emily, S., Ulrika, E., Bo Gullberg, K.S., Hakan, O., and Elisabet, W. (2008). Do both heterocyclic amines and omega-6 polyunsaturated fatty acids contribute to the incidence of breast cancer in postmenopausal women of the Malmo diet and cancer cohort? *The International Journal of Cancer* 123 (7): 163 -1643.
- Friedewald, W .T., R . I . Levy and D . S . Fredrickson, 1972 . Estimation of the concentration of low –density lipoprotein cholesterol in plasma , without use of the preparative ultracentrifuge .*Clin. Chem .*, 18 : 499-502.
- Garg, A., (1994). Efficacy of dietary fiber in lowering serum cholesterol. *Am. J. Med.* 97:501.
- Grootvelt, M., (1998). In vivo absorption, metabolism and urinary excretion of a, b- unsaturated aldehydes in experimental animals: Relevance to the development of cardiovascular diseases by the dietary ingestion of thermally stressed poly unsaturated- rich culinary oils. *J. Clin. Invest.* 101:1210.
- Gurr, M., I., Nazeli, B., and Smith, G. (1989). Dietary fat and plasma lipids. *Nutr. Res. Rev.*, 2:63-86.
- Hajji, T., Khosla P., Pronzkw, A. and Hayes, K. C. (1998). Myristic acid – rich fat raises plasma LDL by stimulating LDL production without affecting fractional clearance in gerbils fed a cholesterol free diet. *J. Nutr.* , 128: 447-454.

- Hibbeln, J.R., Levi, R.G., and Lands. M. (2006). Healthy intakes of n-3 and n-6 fatty acids. *American Journal of Clinical Nutrition* 83 (6):1483S- 1493 S.
- Hodeson, L., Skeaf, C.M. and Chisholm, W. (2001). The effect of replacing dietary saturated fat with poly unsaturated or mono unsaturated fat on plasma lipids in free – living young adults. *Euro. J. Clin. Nutr.* 55:908 – 15.
- Hui, Y. H., (1992). *Encyclopedia of food science and technology*. 1:406 – 16 John Wiley and Sons, Inc. New York.
- Ide, T., Tanaka, T. and Sugano, (1979). Dietary fat dependent changes in hepatic cholesterogenesis and the activity of 3 – hydroxyl – 3-methylglutary – Co A reductase in fasted reefered rats. *J. Nutr.* 109:807 – 18.
- Ishmael, A.L. (1989). Effect of vegetable oils components on oils stability PhD. Thesis, Fac. Of Agric., Cairo Univ., pp. 61:63.
- John, H., Foster, V., Badimon, A. and Chesbro, J. (1990) .Reviews syndromes of accelerate atherosclerosis: role of vascular injury and smooth muscle cell proliferation. *J. Am.Coll.Cardiol.*, 15:1667-87.
- Koo, C., Innera, T. and Mahley, W. (1985). Obligatory role of cholesterol and apolipoprotein-E in the formation of large cholesterol- enriched and receptor- active high density lipoproteins. *J Biol. Chem.* 260:11934 -43.
- Lands, W. E. M. (2005). Dietary fat and health: the evidence and the politics of Prevention: careful use of dietary fats can improve life and prevent disease. *Annals of the New York Academy of Sciences* (Blackwell) 1055: 179- 192. Doi: 10.1196/ annals. 1323.028. PMID 16387724.
- Morrison, W .R. and L . M. Smith , 1962 . Preparation of fatty acid methyl ester and dimethyl acetals from lipid with boron flourid –methanol . *J . lipid . Res .* , 5: 600.
- NCEP, (1993). Summary of the second report of the national cholesterol education program expert panel on detection. Evaluation and treatment of high blood cholesterol in adults. *J. A. M. A.*, 269:3015- 23.
- Okuyama, H., Ichikawa, Y., Sun, Y., Hamazaki, T., and Lands, W. E. M. (2007). Omega-3 fatty acids effectively prevent coronary heart disease and other late- onset diseases: the excessive linoleic acid syndrome. *World Review of Nutritional Dietetics* (Karger) 96 (Prevention of Coronary Heart Disease): 83- 103. PMID 17167282
- Plummer, D. T., (1978). *An introduction to practical biochemistry*, McGraw-Hill Book Company Limited, London.
- Rawashdeh, A. Y. A., (2002). Influences of olive oil and Ghee (samen balady) on serum cholesterol of Jordanians. *Pakistan, J. of Nutr.* 1 (6): 270- 275.
- Rossouw, J. Elza-Mane, B. and Johont, F. (1981). The effect of skim milk, Yoghurt, and full cream milk on human serum lipids. *Am. J. Clin. Nutr.* 34; 351-56.
- Simopoulos, A.P. (2002). Importance of the ratio of omega-6/ omega-3 essential fatty acids. *Biomedicine & Pharmacotherapy.* 56 (8): 365- 379.

- Stahl, E. (1967). Thin layer chromatography. A laboratory Hand book, p.359, springer Verlag, Berlin, New York.
- Valeria, P., Vittorio, K., Paola, M. S., Sabina, S., and Franco, B. (2001). Erythrocyte Membrane Fatty Acids and Subsequent Breast Cancer. JNCL .93 (14):1088. PMID 11459870.
- Visioli, .F.and Galli, C. (1999). The effect of minor constituents of olive oil on cardiovascular disease: new findings. Nutr. Rev., 56:142- 147.
- U .S. Department of Agriculture, Agricultural Research Service. (2007). USDA National Nutrient Database for Standard Reference, Release 20. Nutrient Data Laboratory Home Page, Listed under " Oil, corn industrial and retail, all purpose salad or cooking."
- Yang, Q., Chen, M., and Paola, M. (2007). Modulation of prostate cancer genetic risk by omega-3 and omega-6 fatty acids. The Journal of Clinical Investigation 117 (7): 1866-1875.

### تأثير التغذية البشرية من خلال وجبات تحتوي على السمن وزيت الزيتون ومزيج من هذه الدهون على الكوليسترول في الدم.

- عزت محمد إبراهيم عابدين<sup>١</sup> ، عدلي سمير عبد الستار<sup>٢</sup> ، عماد عبد الفتاح السرسى<sup>٣</sup> ، عبد الستار طلبية جبر<sup>٤</sup> ، محمود عبد الفتاح<sup>١</sup> و عواطف إبراهيم إسماعيل<sup>٢</sup>
- ١- قسم الخبز والعجائن ,معهد بحوث تكنولوجيا الأغذية , مركز البحوث الزراعية, الجيزة , مصر.  
٢- قسم الزيوت , معهد بحوث تكنولوجيا الأغذية , مركز البحوث الزراعية , الجيزة , مصر.  
٣- كلية التربية الرياضية جامعة طنطا.  
٤- قسم الألبان- معهد بحوث تكنولوجيا الأغذية-مركز البحوث الزراعية-الجيزة-مصر.

تم إجراء هذا البحث بغرض عدم توفر كمية الذرة الكافية للحصول على زيت الذرة النقي المتواجد بكمية كبيرة بالأسواق. والسبب أن وزن الجنين يمثل ٤% من وزن حبة الذرة كما أن نسبة الزيت في جنين الذرة ٢٧:٢٥% من وزن الجنين. لذا يلزم طن ذرة للحصول على ١٠ كجم زيت إذا كان الاستخلاص ١٠٠% وهذا مستحيل . كما أظهرت دراسات لبعض الباحثين أن زيت الذرة وزيوت صناعية أخرى تنمي خلايا سرطان البروستاتة وزيادة معدلات الوفاة . كذلك رغبة جمهور المستهلكين على التغذية بالسمن لما لها من مذاق ونكهة مميزة وأثار حميدة على خلايا المخ وتخوفهم في نفس الوقت من ارتفاع نسبة الكوليسترول في الدم . حيث أجريت تجارب تغذية إنسان على وجبات بالسمن البلدي وأخرى بزيت الزيتون كلا على انفراد وخليط منهم .

تم تحليل للأحماض الدهنية المتواجدة بزيت الزيتون والسمن الجاموسى وتم وضع تصميم للتجارب على أساس ثلاثة متغيرات وكل متغير على ثلاثة مستويات كالآتى :

١- ممارسة النشاط الرياضي على مستويات ثلاثة ( لا يمارس أي نشاط – يمارس دون انتظام – يمارس بانتظام ) بواسطة برنامج تدريبي مقترح لتنمية عنصر التوازن أثناء الثبات والحركة في الكاتا الإيجابية لرياضة الكاراتيه . ويتضمن البرنامج أساليب الأعداد البدني والتمارين اللازمة لبناء عضلات الطرف العلوي (الساعد) للكم والطرف السفلي (الفخذ) للركل. والمهارات اللازمة للمجموعة التي تمارس رياضة الكاراتيه بانتظام. وإخضاعهم لبرنامج غذائي بكلية التربية الرياضية – جامعة كفر الشيخ .

٢- الفئة العمرية ومستوياتها ( أقل من ٢٠ سنة – من ٢٠ : ٣٠ سنة – أكثر من ٣٠ سنة ) .

٣- السمنة ومستوياتها ( نحيف – متوسط – سمين ) .

كما تم سحب عينة دم من كل الأفراد المشاركة في التجربة قبل بدء التجارب وتقدير الجلسريدات الثلاثية (TG) والليبيروتين منخفض (LDL-c) وعالي (HDL-c) الكثافة والكولسترول الكلي (TC) واستبعاد العينات الغير مطابقة (Reference range) . أجريت التغذية لمدة ثلاثة أسابيع متتالية ثم تم سحب عينة دم وأخرى بعد التغذية بأسبوع. وحساب المتوسطات لكل من HDL-c, LDL-c, TG , TC . وأتضح الأتي :  
لا يحتوى زيت الزيتون على أحماض دهنية قصيرة السلسلة حتى كربون ١٥ مقارنة بالسمن البلدي الذي يبدأ بكربون ٤ .

وجد أن التغذية بوجبات تحتوى على السمن ترفع نسبة الكولسترول حوالي ٧% وتخفض الجلسريدات الثلاثية . وأن ممارسة أنشطة رياضية والفئة العمرية أقل من ٢٠ سنة والمتوسطة الوزن أقل تأثير من غيرها

كما وجد أن التغذية بوجبات تحتوى على زيت الزيتون تخفض نسبة الكولسترول بما فيها الكولسترول عالي الكثافة وترفع من نسبة الجلسريدات الثلاثية وقد يكون ذلك أيضا خطر على الصحة العامة وقد تلاحظ أن ممارسة النشاط الرياضي يقلل من الجلسريدات الثلاثية ويخفض نسبة الكولسترول وخاصة المنخفض الكثافة

وبالنسبة للتغذية على الوجبات التي تحتوى على خليط من السمن وزيت الزيتون تلاحظ أنه لا يوجد تغير يذكر سيات في مستوى الكولسترول أو الجلسريدات الثلاثية للأفراد التي تزاوول أنشطة رياضية . وعلى هذا الأساس تعتبر التغذية بخلط من السمن وزيت الزيتون تغذية صحية آمنة لتفادى الإصابة بأمراض الأوعية والقلب والشرابين . وخصوصا عند ممارسة النشاط الرياضي لبناء كتل من العضلات والتغلب على السممة الزائدة.

ننصح بدراسة جيدة للزيوت النباتية عموما لبيان النسب بين أوميغا ٦- وأوميغا-٣ وتعديلها عن طريق خلط الزيوت النباتية ببعضها لتصبح النسبة بين أوميغا ٦- : أوميغا-٣ هي ٤ : ١ أي الخلط بغرض تعديل النسب لتفادى الإصابة بالسرطان.

#### قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة  
كلية التربية النوعية – جامعة طنطا

أ.د / مسعد عبد العزيز أبو رية  
أ.د / عادل عبد الحميد الباجورى